

**SUPPLEMENTARY 1:**

Excluded studies from the the workstream 1 literature search strategy, with rationale.

	Authors, year	Reason for exclusion
1.	Alikhan et al., 2004 <sup>1</sup>	Derivation only
2.	Alper et al., 2018 <sup>2</sup>	Not a RAM for predicting the risk of developing VTE in hospital inpatients
3.	Arcelus et al., 1991 <sup>3</sup>	No relevant/useable outcome data
4.	Arpaia et al., 2020 <sup>4</sup>	Not a RAM for predicting the risk of developing VTE in hospital inpatients
5.	Bagot et al., 2010 <sup>5</sup>	No relevant/useable outcome data
6.	Bellizzi et al., 2018 <sup>6</sup>	No relevant/useable outcome data
7.	Blondon et al., 2020 <sup>7</sup>	Duplicate of included study (Blondon et al., 2019a)
8.	Blondon et al., 2017 <sup>8</sup>	Abstract of an included full-text study
9.	Caprini et al., 2001 <sup>9</sup>	Review
10.	Chen et al., 2018 <sup>10</sup>	Not available (foreign language)
11.	Chopard et al., 2006 <sup>11</sup>	No relevant/useable outcome data
12.	Coelho et al., 2020 <sup>12</sup>	Not a RAM for predicting the risk of developing VTE in hospital inpatients
13.	Dang et al., 2019 <sup>13</sup>	Derivation only
14.	Davis and Intagliata, 2018 <sup>14</sup>	No relevant/useable outcome data
15.	Depietri et al., 2018 <sup>15</sup>	No relevant/useable outcome data
16.	Ellis et al., 2019 <sup>16</sup>	No relevant/useable outcome data
17.	Fadoi Foundation, 2020 <sup>17</sup>	Trial protocol (no results)
18.	Ferreira et al., 2018 <sup>18</sup>	No relevant/useable outcome data
19.	Ferreira et al., 2017 <sup>19</sup>	No relevant/useable outcome data
20.	Fritz et al., 2021 <sup>20</sup>	No relevant/useable outcome data
21.	Gibson et al., 2017 <sup>21</sup>	RAM involves diagnostic testing
22.	Girardi et al., 2018 <sup>22</sup>	Not a RAM for predicting the risk of developing VTE in hospital inpatients
23.	Grzelak et al., 2019 <sup>23</sup>	Not a RAM for predicting the risk of developing VTE in hospital inpatients
24.	Hostler et al., 2016 <sup>24</sup>	No relevant/useable outcome data
25.	Hu, 2018 <sup>25</sup>	No relevant/useable outcome data
26.	Koren et al., 2017 <sup>26</sup>	Not a RAM for predicting the risk of developing VTE in hospital inpatients
27.	Kucher et al., 2005 <sup>27</sup>	No relevant/useable outcome data
28.	Lecumberri et al., 2008 <sup>28</sup>	No relevant/useable outcome data
29.	Luo and Zhang, 2017 <sup>29</sup>	Not available (foreign language)

## Supplementary material

30.	Maynard et al., 2010 <sup>30</sup>	No relevant/useable outcome data
31.	McCaffrey et al., 2007 <sup>31</sup>	No relevant/useable outcome data
32.	Meizoso et al., 2017 <sup>32</sup>	Derivation only
33.	Monti et al., 2019 <sup>33</sup>	Abstract of included full text paper
34.	Mull et al., 2017 <sup>34</sup>	Review
35.	Nafee et al., 2018 <sup>35</sup>	Abstract of included full text paper
36.	Nnadi et al., 2017 <sup>36</sup>	No relevant/useable outcome data
37.	Obi et al., 2015 <sup>37</sup>	Critical care patients
38.	Pannucci et al., 2017 <sup>38</sup>	Review
39.	Rafizadeh et al., 2016 <sup>39</sup>	No relevant/useable outcome data
40.	Rastogi et al., 2020 <sup>40</sup>	No relevant/useable outcome data
41.	Razak et al., 2019 <sup>41</sup>	Not a RAM for predicting the risk of developing VTE in hospital inpatients
42.	Robert-Ebadi et al., 2017 <sup>42</sup>	Not a RAM for predicting the risk of developing VTE in hospital inpatients
43.	Salim et al., 2018 <sup>43</sup>	No relevant/useable outcome data
44.	Samama et al., 2006 <sup>44</sup>	Expert opinion
45.	Shrotriya et al., 2018 <sup>45</sup>	Not a RAM for predicting the risk of developing VTE in hospital inpatients
46.	Smilg Nicolas et al., 2018 <sup>46</sup>	No relevant/useable outcome data
47.	Spirk et al., 2017 <sup>47</sup>	No relevant/useable outcome data
48.	Spyropoulos et al., 2020 <sup>48</sup>	RAM involves diagnostic testing
49.	Spyropoulos et al., 2011 <sup>49</sup>	Derivation only
50.	Stuck et al., 2017 <sup>50</sup>	Review
51.	Tadesse et al., 2020 <sup>51</sup>	No relevant/useable outcome data
52.	Taha et al., 2020 <sup>52</sup>	No relevant/useable outcome data
53.	Tung et al., 2020 <sup>53</sup>	No relevant/useable outcome data
54.	Veith et al., 2019 <sup>54</sup>	No relevant/useable outcome data
55.	Winoker et al., 2017 <sup>55</sup>	Abstract of an included full-text study
56.	Yale et al., 2005 <sup>56</sup>	Derivation only
57.	Ye et al., 2017 <sup>57</sup>	Review
58.	Zakai et al., 2013 <sup>58</sup>	Derivation only
59.	Zambelli et al., 2020 <sup>59</sup>	Not a RAM for predicting the risk of developing VTE in hospital inpatients
60.	Zhou et al., 2012 <sup>60</sup>	Not a RAM for predicting the risk of developing VTE in hospital inpatients

## Supplementary material

### References:

1. Alikhan R, Cohen AT, Combe S, et al. Risk Factors for Venous Thromboembolism in Hospitalized Patients With Acute Medical Illness: Analysis of the MEDENOX Study. *JAMA Internal Medicine* 2004;164(9):963-68. doi: 10.1001/archinte.164.9.963
2. Alper EC, Ip IK, Balthazar P, et al. Risk Stratification Model: Lower-Extremity Ultrasonography for Hospitalized Patients with Suspected Deep Vein Thrombosis. *Journal of General Internal Medicine* 2018;33(1):21-25. doi: <https://dx.doi.org/10.1007/s11606-017-4170-3>
3. Arcelus JI, Candocia S, Traverso CI, et al. Venous thromboembolism prophylaxis and risk assessment in medical patients. *Semin Thromb Hemost* 1991;17:3313-8.
4. Arpaia GG, Caleffi A, Marano G, et al. Padua prediction score and IMPROVE score do predict in-hospital mortality in Internal Medicine patients. *Internal and Emergency Medicine* 2020;15(6):997-1003. doi: <https://dx.doi.org/10.1007/s11739-019-02264-4>
5. Bagot C, Gohil S, Perrott R, et al. The use of an exclusion-based risk-assessment model for venous thrombosis improves uptake of appropriate thromboprophylaxis in hospitalized medical patients. *QJM* 2010;103:597-605.
6. Bellizzi A, Palombi G, Bardi G, et al. Score Padua and IMPROVE: What changes in management of venous thromboembolism prophylaxis in Internal Medicine? *Italian Journal of Medicine* 2018;12(2 Supplement 1):2. doi: <http://dx.doi.org/10.4081/itjm.2018.s2>
7. Blondon M, Righini M, Nendaz M, et al. External validation of the simplified Geneva risk assessment model for hospital-associated venous thromboembolism in the Padua cohort. *Journal of Thrombosis and Haemostasis* 2020;18(3):676-80. doi: <https://dx.doi.org/10.1111/jth.14688>
8. Blondon M, Spirk D, Kucher N, et al. External validation and comparison of the improve risk assessment model with the Geneva risk assessment model in the estimate cohort. *Research and Practice in Thrombosis and Haemostasis* 2017;1(Supplement 1):180. doi: <http://dx.doi.org/10.1002/rth2.12012>
9. Caprini JA, Arcelus JI, Reyna J. Effective risk stratification of surgical and nonsurgical patients for venous thromboembolic disease. *Seminars in Hematology* 2001;38:12-19. doi: [https://doi.org/10.1016/S0037-1963\(01\)90094-0](https://doi.org/10.1016/S0037-1963(01)90094-0)
10. Chen XL, Pan L, Wang Y. [Validity of Padua risk assessment scale for assessing the risk of deep venous thrombosis in hospitalized patients]. *Zhonghua nei ke za zhi* 2018;57(7):514-17. doi: <https://dx.doi.org/10.3760/cma.j.issn.0578-1426.2018.07.009>
11. Chopard P, Spirk D, Bounameaux H, et al. Identifying acutely ill medical patients requiring thromboprophylaxis. *J Thromb Haemost* 2006;4(4):915-6.
12. Coelho J, Divernet-Queriaud M, Roy P-M, et al. Comparison of the Wells score and the revised Geneva score as a tool to predict pulmonary embolism in outpatients over age 65. *Thrombosis Research* 2020;196:120-26. doi: <https://dx.doi.org/10.1016/j.thromres.2020.07.026>
13. Dang JT, Switzer N, Delisle M, et al. Predicting venous thromboembolism following laparoscopic bariatric surgery: development of the BariClot tool using the MBSAQIP database. *Surgical Endoscopy* 2019;33(3):821-31. doi: <https://dx.doi.org/10.1007/s00464-018-6348-0>
14. Davis J, Intagliata NM. Practice patterns in application of venous thromboembolism prophylaxis to a hepatology inpatient service: A quality improvement project. *Hepatology* 2018;68(Supplement 1):296A. doi: <http://dx.doi.org/10.1002/hep.30257>

## Supplementary material

15. Depietri L, Marietta M, Scarlini S, et al. Clinical impact of application of risk assessment models (Padua Prediction Score and Improve Bleeding Score) on venous thromboembolism, major hemorrhage and health expenditure associated with pharmacologic VTE prophylaxis: a "real life" prospective and retrospective observational study on patients hospitalized in a Single Internal Medicine Unit (the STIME study). *Internal and Emergency Medicine* 2018;13(4):527-34. doi: <https://dx.doi.org/10.1007/s11739-018-1808-z>
16. Ellis N, Grubb C-M, Mustoe S, et al. Venous thromboembolism risk in psychiatric inpatients: a multicentre cross-sectional study. *BJPsych Bulletin* 2019;1-5. doi: <https://dx.doi.org/10.1192/bjb.2019.25>
17. Fadot Foundation. Efficacy of the Use of Risk Scores in Reducing Important Clinical Outcomes in Hospitalized Medical III Patients, 2020. Available <https://clinicaltrialsgov/show/NCT04267718> [Accessed 2 March 2021]
18. Ferreira C, Avelino Mancini R, Monique Lana Diniz M, et al. Evaluation of inter-rater agreement of a risk assessment model for thromboprophylaxis in medical hospitalised patients: Preliminary results of a cohort study. *Research and Practice in Thrombosis and Haemostasis* 2018;2(Supplement 1):323-24. doi: <http://dx.doi.org/10.1002/rth2.12125>
19. Ferreira D, de Sousa JA, Felicissimo P, et al. Venous thromboembolism risk and prophylaxis in the Portuguese hospital care setting: The ARTE study. *Revista portuguesa de cardiologia : orgao oficial da Sociedade Portuguesa de Cardiologia (Portuguese Journal of Cardiology)* 2017;36(11):823-30. doi: <https://dx.doi.org/10.1016/j.repc.2017.01.009>
20. Fritz MK, Kincaid SE, Sargent CG, et al. Venous thromboembolism (VTE) risk stratification in general medical patients at an academic medical center. *Journal of Thrombosis and Thrombolysis* 2021;51(1):67-73. doi: <https://dx.doi.org/10.1007/s11239-020-02144-1>
21. Gibson CM, Spyropoulos AC, Cohen AT, et al. The IMPROVED VTE Risk Score: Incorporation of D-Dimer into the IMPROVE Score to Improve Venous Thromboembolism Risk Stratification. *TH Open* 2017;1(1):e56-e65. doi: <https://dx.doi.org/10.1055/s-0037-1603929>
22. Girardi AM, Bettoli RS, Garcia TS, et al. Wells and Geneva Scores Are Not Reliable Predictors of Pulmonary Embolism in Critically Ill Patients: A Retrospective Study. *Journal of Intensive Care Medicine* 2018;885066618816280. doi: <https://dx.doi.org/10.1177/0885066618816280>
23. Grzelak M, Javia V, Cox C, et al. Incidence and patient characteristics for the development of venous thromboembolic events within the adult burn center population and an evaluation of the caprini risk assessment model. *Journal of Burn Care and Research* 2019;40(Supplement 1):S57. doi: <http://dx.doi.org/10.1093/jbcr/irz013.088>
24. Hostler DC, Marx ES, Moores LK, et al. Validation of the international medical prevention registry on venous thromboembolism bleeding risk score. *Chest* 2016;149(2):372-79. doi: <http://dx.doi.org/10.1378/chest.14-2842>
25. Hu Y. Comparison between caprini and Khorana risk assessment models for hospitalized medical patients with cancer at high risk for venous thromboembolism: a retrospective study. *American Journal of Respiratory and Critical Care Medicine* 2018;197(MeetingAbstracts)
26. Koren O, Hourri A, Elias M, et al. Failure to validate padua score's ability to predict the risk of venous thromboembolism in medical patients. *Clinical Therapeutics* 2017;39(8 Supplement 1):e59.
27. Kucher N, Koo S, Quiroz R, et al. Electronic alerts to prevent venous thromboembolism among hospitalized patients. *N Engl J Med* 2005;352(10):969-77.

## Supplementary material

28. Lecumberri R, Marqués M, Díaz-Navarla MT, et al. Maintained effectiveness of an electronic alert system to prevent venous thromboembolism among hospitalized patients. *Thromb Haemost* 2008;100(4):699-704.
29. Luo X, Zhang F. Validation of the Caprini risk assessment model for venous thromboembolism in Chinese hospitalized patients in a general hospital. *National Medical Journal of China* 2017;97(24):1875-77. doi: <http://dx.doi.org/10.3760/cma.j.issn.0376-2491.2017.24.007>
30. Maynard GA, Morris TA, Jenkins IH, et al. Optimizing prevention of hospital-acquired venous thromboembolism (VTE): prospective validation of a VTE risk assessment model. *J Hosp Med* 2010;5(1):10-18.
31. McCaffrey R, Bishop M, Adonis-Rizzo M, et al. Development and Testing of a DVT Risk Assessment Tool: Providing Evidence of Validity and Reliability. *Worldviews on Evidence-Based Nursing* 2007;4(1):14-20. doi: 10.1111/j.1741-6787.2007.00074.x
32. Meizoso JP, Karcutskie CA, Ray JJ, et al. A simplified stratification system for venous thromboembolism risk in severely injured trauma patients. *The Journal of Surgical Research* 2017;207:138-44. doi: <https://dx.doi.org/10.1016/j.jss.2016.08.072>
33. Monti M, Timpone S, Murdolo G, et al. A new risk assessment model for the stratification of the thromboembolism risk in medical patients: The TEVere Score. *Italian Journal of Medicine* 2019;13(Supplement 2):7-8. doi: <http://dx.doi.org/10.4081/itjm.2019.s2>
34. Mull N, Hecht T, Umscheid C, et al. Clinical risk prediction scores for venous thromboembolism in hospitalized patients. *International Journal of Technology Assessment in Health Care* 2017;33(Supplement 1):75-76. doi: <http://dx.doi.org/10.1017/S0266462317002094>
35. Nafee T, Gibson CM, Travis R, et al. Performance of a machine learning model vs. IMPROVE score for VTE prediction in acute medically ill patients: Insights from the APEX trial. *European Heart Journal* 2018;39(Supplement 1):431. doi: <http://dx.doi.org/10.1093/eurheartj/ehy565.2160>
36. Nnadi C, Soltani SJ, Brice HG, et al. Evaluation of a new thrombo-embolism scoring system (TESS) for spinal surgery. *Spine Journal* 2017;17(3):S21-S22. doi: <http://dx.doi.org/10.1016/j.spinee.2016.12.068>
37. Obi AT, Pannucci CJ, Nackashi A, et al. Validation of the Caprini Venous Thromboembolism Risk Assessment Model in Critically Ill Surgical Patients. *JAMA Surgery* 2015;150(10):941-48. doi: 10.1001/jamasurg.2015.1841
38. Pannucci CJ, Swistun L, MacDonald JK, et al. Individualized Venous Thromboembolism Risk Stratification Using the 2005 Caprini Score to Identify the Benefits and Harms of Chemoprophylaxis in Surgical Patients: A Meta-analysis. *Annals of Surgery* 2017;265(6):1094-103. doi: <https://dx.doi.org/10.1097/SLA.0000000000002126>
39. Rafizadeh R, Turgeon RD, Batterink J, et al. Characterization of venous thromboembolism risk in medical inpatients using different clinical risk assessment models. *Canadian Journal of Hospital Pharmacy* 2016;69(6):454-59.
40. Rastogi R, Mehaffey JH, Turrentine BE, et al. Implementation of an Electronic Medical Record Risk Stratification Tool to Reduce Venous Thromboembolism Events. *Journal of the American College of Surgeons* 2020;231(4 Supplement 1):S141-S42. doi: <http://dx.doi.org/10.1016/j.jamcollsurg.2020.07.273>
41. Razak MKA, Rasheed JI, Hamad AA. A comparison between wells' score and modified doppler ultrasound in the diagnosis of deep venous thrombosis of the lower limb.

## Supplementary material

- International Journal of Pharmaceutical Sciences and Research* 2019;10(5):2369-74. doi: <http://dx.doi.org/10.13040/IJPSR.0975-8232.10%285%29.2369-74>
42. Robert-Ebadi H, Mostaguir K, Hovens MM, et al. Assessing clinical probability of pulmonary embolism: prospective validation of the simplified Geneva score. *Journal of Thrombosis and Haemostasis* 2017;15(9):1764-69. doi: <https://dx.doi.org/10.1111/jth.13770>
43. Salim S, Onida S, Davies A. Variability in venous thromboembolism (VTE) risk assessment tools. *Phlebology* 2018;33(1 Supplement 1):36-37. doi: <http://dx.doi.org/10.1177/0268355518817748>
44. Samama MM, Dahl OE, Mismetti P, et al. An electronic tool for venous thromboembolism prevention in medical and surgical patients. *Haematologica* 2006;91(1):64.
45. Shrotriya S, Dhakal P, Sharma M, et al. Predictors of venous thromboembolism among cancer inpatients: A predictive model. *Journal of Clinical Oncology* 2018;36(15 Supplement 1) doi: [http://dx.doi.org/10.1200/JCO.2018.36.15\\_suppl.e18824](http://dx.doi.org/10.1200/JCO.2018.36.15_suppl.e18824)
46. Smilg Nicolas C, Tornel Sanchez G, Trujillo Santos J. Concordance among venous thromboembolism risk assessment models. *Concordancia entre las escalas de predicción de riesgo de tromboembolia venosa* 2018;150(2):61-63. doi: <https://dx.doi.org/10.1016/j.medcli.2017.06.021>
47. Spirk D, Stuck AK, Hager A, et al. Electronic alert system for improving appropriate thromboprophylaxis in hospitalized medical patients: a randomized controlled trial. *Journal of Thrombosis and Haemostasis* 2017;15(11):2138-46. doi: <https://dx.doi.org/10.1111/jth.13812>
48. Spyropoulos AC, Lipardi C, Xu J, et al. Modified IMPROVE VTE Risk Score and Elevated D-Dimer Identify a High Venous Thromboembolism Risk in Acutely Ill Medical Population for Extended Thromboprophylaxis. *TH Open* 2020;4(1):e59-e65. doi: <https://dx.doi.org/10.1055/s-0040-1705137>
49. Spyropoulos AC, Anderson FA, Fitzgerald G, et al. Predictive and Associative Models to Identify Hospitalized Medical Patients at Risk for VTE. *Chest* 2011;140(3):706-14. doi: <https://doi.org/10.1378/chest.10-1944>
50. Stuck AK, Spirk D, Schaudt J, et al. Risk assessment models for venous thromboembolism in acutely ill medical patients. A systematic review. *Thrombosis and Haemostasis* 2017;117(4):801-08. doi: <https://dx.doi.org/10.1160/TH16-08-0631>
51. Tadesse TA, Kedir HM, Fentie AM, et al. Venous Thromboembolism Risk and Thromboprophylaxis Assessment in Surgical Patients Based on Caprini Risk Assessment Model. *Risk Management and Healthcare Policy* 2020;13:2545-52. doi: <https://dx.doi.org/10.2147/RMHP.S272852>
52. Taha H, Govindraj E, Jaber F, et al. Improving venous thromboembolism prophylaxis through critical thinking and health informatics. *BMJ Open Quality* 2020;9(2) doi: <https://dx.doi.org/10.1136/bmjoq-2019-000885>
53. Tung EC, Yu S-Y, Shah K, et al. Reassessment of venous thromboembolism and bleeding risk in medical patients receiving VTE prophylaxis. *Journal of Evaluation in Clinical Practice* 2020;26(1):18-25. doi: <https://dx.doi.org/10.1111/jep.13213>
54. Veith J, Collier W, Rockwell WB, et al. Direct Comparison of Patient-completed and Physician-completed Caprini Scores for Plastic Surgery Patients. *Plastic and Reconstructive Surgery Global Open* 2019;7(8):e2363. doi: <https://dx.doi.org/10.1097/GOX.0000000000002363>

## Supplementary material

55. Winoker JS, Paulucci DJ, Anastos H, et al. MP76-07 Predicting complications following robot-assisted partial nephrectomy with the ACS-NSQIP universal surgical risk calculator. *Journal of Urology* 2017;197(4S):e1015.
56. Yale S, Medlin S, Liang H, et al. Risk assessment model for venothromboembolism in post-hospitalized patients. *International Angiology* 2005;24(3):250.
57. Ye F, Stalvey C, Khuddus MA, et al. A systematic review of mobility/immobility in thromboembolism risk assessment models for hospitalized patients. *Journal of Thrombosis and Thrombolysis* 2017;44(1):94-103. doi: <https://dx.doi.org/10.1007/s11239-017-1501-5>
58. Zakai NA, Callas PW, Repp AB, et al. Venous thrombosis risk assessment in medical inpatients: the medical inpatients and thrombosis (MITH) study. *J Thromb Haemost* 2013;11(4):634-41.
59. Zambelli R, Nemeth B, Touw C, et al. Is the caprini score a reliable tool to predict the risk of venous thrombosis in orthopedic patients. *Research and Practice in Thrombosis and Haemostasis* 2020;4(SUPPL 1):1194-95. doi: <http://dx.doi.org/10.1002/rth2.12393>
60. Zhou H-X, Peng L-Q, Yan Y, et al. Validation of the Caprini risk assessment model in Chinese hospitalized patients with venous thromboembolism. *Thrombosis Research* 2012;130(5):735-40. doi: <https://doi.org/10.1016/j.thromres.2012.08.001>

## Supplementary material