

Table S7. Baseline characteristics in patients undergoing OPCAB converted vs non-converted to ONCAB

| | OPCAB not converted | OPCAB converted to on-pump | P* |
|-------------------------------|---------------------|----------------------------------|-------|
| n | 1231 | 29 | |
| Age (years) mean (sd) | 63.59 (9.06) | 64.40 (9.77) | 0.634 |
| Female n(%) | 179 (14.5) | 1 (3.4) | 0.16 |
| BMI mean (sd) | 28.10 (4.11) | 28.15 (3.99) | 0.95 |
| SBP (mmHg) mean (sd) | 132.75 (19.11) | 130.66 (17.94) | 0.56 |
| DBP (mmHg) mean (sd) | 75.40 (11.07) | 72.48 (11.85) | 0.16 |
| Creatinine (mmol/l) mean (sd) | 99.68 (22.19) | 107.76 (41.54) | 0.06 |
| Treated Hypertension n(%) | 919 (74.7) | 24 (82.8) | 0.44 |
| Treated Hyperlipaemia n(%) | 1147 (93.2) | 29 (100.0) | 0.28 |
| Diabetes n(%) | | | 0.78 |
| No | 959 (77.9) | 21 (72.4) | |

| | | | |
|-----------------------|------------|------------|-------|
| Insulin dependent | 68 (5.5) | 2 (6.9) | |
| Non-insulin dependent | 204 (16.6) | 6 (20.7) | |
| Smoking n(%) | | | 0.02 |
| Current | 173 (14.1) | 7 (24.1) | |
| Ex-smoker | 645 (52.4) | 19 (65.5) | |
| Never | 413 (33.5) | 3 (10.3) | |
| COPD n(%) | 29 (2.4) | 0 (0.0) | 0.83 |
| Asthma n(%) | 58 (4.7) | 5 (17.2) | 0.009 |
| PVD n(%) | 87 (7.1) | 3 (10.3) | 0.76 |
| TIA n(%) | 40 (3.2) | 0 (0.0) | 0.65 |
| CVA n(%) | 40 (3.2) | 0 (0.0) | 0.65 |
| MI n(%) | 493 (40.0) | 17 (58.6) | 0.07 |
| PCI n(%) | 199 (16.2) | 9 (31.0) | 0.06 |
| Preop AF n(%) | 18 (1.5) | 1 (3.4) | 0.92 |
| preop LVEF (%) | | | 0.37 |

| | | | |
|--------------------|------------|------------|------|
| $\geq 50\%$ (good) | 927 (75.3) | 23 (79.3) | |
| 31-49% (moderate) | 264 (21.4) | 4 (13.8) | |
| $\leq 30\%$ (poor) | 40 (3.2) | 2 (6.9) | |
| LMD n(%) | 276 (22.4) | 6 (20.7) | 1 |
| RA n(%) | 231 (18.8) | 9 (31.0) | 0.15 |
| SVG n(%) | 916 (74.4) | 20 (69.0) | 0.65 |
| BITA n(%) | 582 (47.3) | 16 (55.2) | 0.51 |

*Chi test or t-test

OPCAB: Off-pump coronary artery bypass; ONCAB: On-pump coronary artery bypass; Systolic blood pressure; DPB: Diastolic blood pressure; COPD: Chronic obstructive pulmonary disease; PVD: Peripheral vascular disease; TIA: Transient ischemic attack; CVA: Cerebrovascular accident; MI: Myocardial infarction; PCI: Percutaneous coronary intervention; AF: Atrial fibrillation; LVEF: Left ventricular ejection fraction; LMD: Left main disease; RA: Radial artery BITA: Bilateral internal thoracic arteries, SVG: Saphenous vein graft. Reproduced from: Benedetto U, et al. Off-pump versus on-pump coronary artery bypass grafting: Insights from the Arterial Revascularization Trial. J Thorac Cardiovasc Surg. 2018;155(4):1545-1553.e7. doi:10.1016/j.jtcvs.2017.10.135.