

#### Supplementary material 4

Table showing estimated intervention effects on self-reported quit attempts of at least 24 hours in the previous 3 or 6 months (reported at 3 and 9 months, respectively)

	Number of participants in analysis		Number of outcomes per group		Unadjusted analysis			Adjusted analysis <sup>a</sup>		
	SAU	Intervention	SAU n (%)	Intervention n (%)	Odds ratio <sup>b</sup> (95% CI) p-value	Absolute between group differences in risk <sup>c</sup> (95% CI) p-value	Relative risk <sup>d</sup> (95% CI) p-value	Odds ratio <sup>b</sup> (95% CI) p-value	Absolute between group differences in risk <sup>c</sup> (95% CI) p-value	Relative risk <sup>d</sup> (95% CI) p-value
Quit attempt made in first 3 months	456	456	37 (8.1)	54 (11.8)	1.52 (0.98, 2.36), 0.062	0.04 (-0.00, 0.08), 0.060	1.46 (0.98, 2.17), 0.062	1.53 (0.99, 2.39), 0.058	0.04 (-0.00, 0.08), 0.056	1.47 (0.99, 2.18), 0.058
Quit attempt made between 3 months and 9 months	451	450	68 (15.1)	76 (16.9)	1.14 (0.80, 1.64), 0.458	0.02 (-0.03, 0.07) 0.458	1.12 (0.83, 1.51), 0.459	1.15 (0.80, 1.64), 0.447	0.02 (-0.03, 0.07) 0.447	1.12 (0.83, 1.52) 0.447

<sup>a</sup>Analysis adjusted for stratification variables (HSI and site)

<sup>b</sup> Odds in Intervention group relative to SAU group

<sup>c</sup> The difference in risk subtracting SAU from Intervention

<sup>d</sup> Risk in Intervention group relative to SAU group

Denominator based on Russell Standards (same denominator as used for primary outcome); assumes missing response = no quit attempt made.