

## S6 Bibliography

### Wrong language

Afnan Hamed-Agbariah and Rosenfeld Y . (2015). [the Added Value of Art Therapy for Mothers with Post-Partum Depression in Arabic Society in Israel]. *Harefuah*, 154(9), pp.568-72, 608.

Andaroon N, Kordi M and Kimiaee S A; Esmaily H ;. (2018). Effect of individual counseling program by a midwife on anxiety during pregnancy in nulliparous women. [Persian]. *Iranian Journal of Obstetrics and Gynecology and Infertility*, 20(12), pp.86-95.

Azadi F, Shariat M and Niromanesh S ; Shirazi M ;. (2016). Effectiveness of stress management training on stress reduction in pregnant women. [Persian]. *Tehran University Medical Journal*, 74(7), pp.493-499.

Benhajjoub S, Ladenburger A and Lighezzolo J ; de Tychev ; C ;. (2008). Maternal and perinatal depression and prevention: A clinical and projective approach. [French]. *Evolution Psychiatrique*, 73(2), pp.331-352.

Bowen A, Baetz M and McKee N ; Klebaum N ;. (2008). Optimizing maternal mental health within a primary health care centre: a model program. *Canadian Journal of Community Mental Health*, 27(2), pp.105-116.

Castanon S C and Pinto L J;. (2008). [Use of the Edinburgh Postnatal Depression Scale to detect postpartum depression]. *Revista Medica de Chile*, 136(7), pp.851-8.

Chabrol H, Teissedre F and Santrisse K ; Armitage J ; Saint-Jean M ;. (2001). [Acceptability of antidepressants and psychotherapies in the depressions of the post-partum: inquiry to 198 delivered]. *Encephale*, 27(4), pp.381-2.

Chabrol H, Teissedre F and Saint-Jean M ; Teisseyre N ; Sistac C ; Michaud C ; Roge B ;. (2002). Detection, prevention and treatment of postpartum depression: A randomized, controlled study on a sample of 859 women. *L'Encephale: Revue de psychiatrie clinique biologique et therapeutique*, 28(1), pp.65-70.

Chang M Y and Chen C H;. (2008). [Evidence-based management of perinatal depression]. *Hu Li Tsa Chih - Journal of Nursing*, 55(2), pp.5-9.

Cheng H Y and Huang T Y; Chien L Y; Cheng Y F; Chen F J;. (2016). [The Effects of a Mobile Application Social Support Program on Postpartum Perceived Stress and Depression]. *Hu Li Tsa Chih - Journal of Nursing*, 63(6), pp.52-60.

Chrzan-Detkos M and Kalita L . (2019). The role of early psychological intervention in the prevention and treatment of postpartum depression. [Polish]. *Psychoterapia*, 188(1), pp.47-61.

Collado Maria Assumpta Ortiz, Hatem Marie and Tourne Claude-Emile ;. (2015). Can we explain the depression of mothers who don't have a psychiatric history by the response to inhibition of the action?. *Revue Quebecoise de Psychologie*, 36(1), pp.187-214.

de Noose , Lisa and Garnier Salome ; Richelle Jacqueline ; Lighezzolo-Alnot Joelle ; Hendrick Stephan ; de Tychev ; Claude ;. (2011). Prenatal depression and therapeutic

engagement: A clinical and projective approach. *Psychologie Clinique et Projective*, 17, pp.159-188.

de Sousa Schwengber, Daniela Delias and Alfaya Cristiane ; de Cassia Sobreira Lopes; Rita ; Piccinini Cesar Augusto;. (2003). Interactional orientation as an alternative of precocious parents-baby intervention in the context of maternal depression: Some initial reflections. *Psico*, 34(2), pp.297-316.

Des Rivieres-Pigeon , C and Seguin L ; Brodeur J M; Perreault M ; Boyer G ; Colin C ; Goulet L ;. (2000). [The Edinburgh Postnatal Depression Scale: the validity of its Quebec version for a population low socioeconomic status mothers]. *Canadian Journal of Community Mental Health*, 19(1), pp.201-14.

Fakhri A, Morshedi H and Mohammadi Zeidi ; I ;. (2017). Effect of education based on health belief model with relaxation on anxiety of nulliparouse women. [Persian]. *Scientific Journal of Kurdistan University of Medical Sciences*, 22(2), pp.32-47.

Fang Hui-Chen, Liu Chieh-Yu and Day Hwey-Lian ; Chen Chie-Pein ; Gau Meei-Ling ;. (2011). Uncertainty, stress, and birth outcomes in non-hospitalized, high-risk pregnancy women: the effectiveness of health consultation. *Journal of Nursing & Healthcare Research*, 7(1), pp.13-13.

Glasser S and Barell V . (1999). [Depression scale for research in and identification of postpartum depression]. *Harefuah*, 136(10), pp.764-8, 844, 843.

Hofecker-Fallahpour M, Zinkernagel-Burri Ch and Stockli B ; Wusten G ; Stieglitz R D; Riecher-Rossler A ;. (2003). [Group therapy for depression during early motherhood: first results of a pilot study]. *Nervenarzt*, 74(9), pp.767-74.

Hohm E, Trautmann-Villalba P and Schwarz M ; Hornstein C ;. (2008). In-patient mother baby therapy for postpartum mental disorders - A contribution to prevention in infancy. [German]. *Nervenheilkunde*, 27(6), pp.506-512.

Hok V, Hoisnard G and Simon-Vernier E ; Hauchecorne A ; Thomas A ; Menard C ; Tubach F ; Saïas T ;. (2011). Psychologists at home: Practices, models and issues of a preventive intervention. *Pratiques Psychologiques*, 17(2), pp.119-135.

Hornstein C, Hohm E and Keppler S ; Trautmann-Villalba P ; Wild E ; Schulz R ;. (2009). Prevention of postnatal disorders starts during pregnancy! multi-professional network of expertise to encourage mentally burdened and teenage mothers and their children after birth: "perinatal prevention network in the rhine-neckar region". [German]. *Journal fur Neurologie and Neurochirurgie und Psychiatrie*, 10(1), pp.48-52.

Hribersek D. (2014). Choice of psychopharmaceutical among pregnant women and nursing mothers. [Slovene]. *Farmaceutski Vestnik*, 65(1), pp.31-38.

Huber G and Seelbach-Gobel B . (2009). Postpartum depression should obstetric hospitals perform routine screening?. [German]. *Geburtshilfe und Frauenheilkunde*, 69(12), pp.1097-1100.

Jalenques I and Legrand G . (2009). [Post-partum blues and depression]. *Revue du Praticien*, 59(4), pp.499-506.

- Kapfhammer H P and Klier C M;. (2009). Bipolar affective disorders in pregnancy and puerperium. New developments in diagnostics and treatment. [German]. *Journal fur Neurologie and Neurochirurgie und Psychiatrie*, 10(1), pp.32-36.
- Karamoozian Mahsa, Askarizadeh Ghasem and Darekordi Ali ;. (2015). The impact of prenatal group stress management therapy on anxiety in mothers and newborns' physiological parameters. *Developmental Psychology: Journal of Iranian Psychologists*, 11(43), pp.259-272.
- Kim M H and Ji E S;. (2016). [Effects of a Hospital Based Follow-Up Program for Mothers with Very Low Birth Weight Infants]. *Journal of Korean Academy of Nursing*, 46(1), pp.79-89.
- Kim E M and Hong S. (2018). [Impact of Uncertainty on the Anxiety of Hospitalized Pregnant Women Diagnosed with Preterm Labor: Focusing on Mediating Effect of Uncertainty Appraisal and Coping Style]. *Journal of Korean Academy of Nursing*, 48(4), pp.485-496.
- Komminaho M. (2003). Postpartum depression screening at Rauma Health District mother and baby clinics. *Sairaanhoitaja*, 76(4), pp.23-25.
- Lara Ma, A and Navarro C ; Navarrete L ; Cabrera A ; Almanza J ; Morales F ; Juarez F ;. (2006). Depressive symptoms in pregnancy and associated factors in patients of three health institutions in Mexico City. [Spanish]. *Salud Mental*, 29(4), pp.55-62.
- Lefebvre-Chanson C, Boissou-Bonnet C and Rolland A C;. (2017). Treating postnatal depression with group therapy. *Neuropsychiatrie de l'Enfance et de l'Adolescence*, 65(1), pp.14-20.
- Luis M A, de Oliveira and E R ;. (1998). [Mental disorders during pregnancy, childbirth and puerperium in the region of Ribeiro Preto-SP, Brazil]. *Revista Da Escola de Enfermagem Da Usp*, 32(4), pp.314-24.
- Martinez P, Rojas G and Magana I ;. (2015). Ethnographic analysis at general practice waiting rooms: Implications for postpartum depression detection and treatment. [Spanish]. *Terapia Psicologica*, 33(2), pp.59-68.
- Mirabella F, Michielin P and Piacentini D ; Veltro F ; Barbano G ; Cattaneo M ; Palumbo G ; Gigantesco A ;. (2016). [Effectiveness of a postnatal psychological treatment for women who had screened positive for depression]. *Rivista di Psichiatria*, 51(6), pp.260-269.
- Moschner S L, Achtergarde S and Ramsauer B ;. (2018). [Treatment Satisfaction of Mothers with Postpartum Depression Concerning Circle of Security Intervention]. *Praxis der Kinderpsychologie und Kinderpsychiatrie*, 67(4), pp.351-366.
- Piacentini Daniele, Leveni Daniela and Primerano Giuseppe ; Cattaneo Marina ; Biffi Giuseppe ; Mirabella Fiorino ; Gigantesco Antonella ;. (2009). Results of a manualised cognitive behavioural treatment in postnatal depression. *Psicoterapia Cognitiva e Comportamentale*, 15(3), pp.379-389.
- Piacentini D, Mirabella F and Leveni D ; Primerano G ; Cattaneo M ; Biffi G ; Rozzoni C ; Arcaini F ; Gigantesco A ;. (2011). [Effectiveness of a manualized cognitive-behavioural intervention for postnatal depression]. *Rivista di Psichiatria*, 46(3), pp.187-94.

Pukas A, Suda W and Kwiatkowska-Gruca M ; Behrendt J ; Godula-Stuglik U ;. (2010). Level of anxiety in parents of neonates treated in the NICU and support from family and medical staff. [Polish]. *Pediatrics Polska*, 85(4), pp.341-344.

Radonic E, Hanzevacki M and Jelavic S ; Jurkovic I ; Malekovic H ; Radonic E N; Vrabec B ;. (2018). Improvement of non-institutional treatment and care for women with postpartum depression. [Croatian]. *Acta Medica Croatica*, 72(2), pp.241-246.

Reyhani T, Pour Z S and Heidarzadeh M ; Mousavi S M; Mazloom S R;. (2014). Investigating the effects of spiritual self-care training on psychological stress of mothers with preterm infants admitted in neonatal intensive care unit. [Persian]. *Iranian Journal of Obstetrics and Gynecology and Infertility*, 17(97), pp.18-27.

Rincon-Pabon D and Ramirez-Velez R . (2014). [Postnatal depression in Colombian women: secondary analysis of the 2010 Colombian Demographic and Health Survey]. *Revista de Salud Publica*, 16(4), pp.534-46.

Rojas G, Santelices M P and Martinez P ; Tomicic A ; Reinel M ; Olhaberry M ; Krause M ;. (2015). [Barriers restricting postpartum depression treatment in Chile]. *Revista Medica de Chile*, 143(4), pp.424-32.

Rojas G, Guajardo V and Martinez P ; Fritsch R ;. (2018). [Screening and barriers for treatment of postpartum depression in Chilean public primary health care centers]. *Revista Medica de Chile*, 146(9), pp.1001-1007.

Saias T, David M and Guedeney A ; Tereno S ; Tubach F ; Greacen T ; Dugravier R ;. (2016). Prevention of perinatal depression in high-risk families through home visitation: Clinical issues. *Pratiques Psychologiques*, 22(2), pp.159-175.

Tang H and Huang Q . (2019). [Comparative study of minimally invasive technique and traditional decaying method for dental caries and the impact of dental treatment on pain and anxiety during pregnancy]. *Shanghai Kou Qiang Yi Xue/Shanghai Journal of Stomatology*, 28(3), pp.297-300.

Varo Inmaculada Suárez-Varela. (2012). Impact of a nursing intervention on pregnant women as a preventive tool for postpartum depression. *Nure Investigación*, (60), pp.1-17.

Yamashita H and Yoshida K . (2003). [Screening and intervention for depressive mothers of new-born infants]. *Seishin Shinkeigaku Zasshi - Psychiatria et Neurologia Japonica*, 105(9), pp.1129-35.

Yin Y H and Wang M. (2017). Impact of nucleos(t)ide analogue antiviral therapy on anxiety and depression in pregnant women with chronic hepatitis B. [Chinese]. *World Chinese Journal of Digestology*, 25(26), pp.2360-2365.

Zelkowitz P and Milet T H. (1995). Screening for post-partum depression in a community sample. *Canadian Journal of Psychiatry - Revue Canadienne de Psychiatrie*, 40(2), pp.80-6.

### **Full text not available**

Aigbe Aitebureme Oare. (2015). *Providers' perspectives on barriers to care, system constraints and piloting the HIV-modified centering pregnancy model among HIV-positive pregnant women*. . Aigbe, Aitebureme Oare: U Texas School of Public Health, US.

Bhati Sulochani Rao. (2015). *The effect of the Sleep Support for Moms Intervention on postpartum sleep and depressive symptoms. a pilot Randomized Controlled Trial.* . Bhati, Sulochani Rao: George Mason U., US.

Brown H and Bacigalupo R . (2006). Health visitors and postnatal depression: identification and practice. *Community Practitioner*, 79(2), pp.49-52.

Brown Michelle D. (2018). The use of mobile technology in managing mild depressive symptomology in the Primipara of low socioeconomic status: A feasibility study. *Dissertation Abstracts International: Section B: The Sciences and Engineering*, 79(9-B(E)), pp.No Pagination Specified.

Cubison Jan and Munro Jane . (2005). Acceptability of Using the EPDS as a Screening Tool for Postnatal Depression. In: , ed., *Screening for perinatal depression*. London, England: Jessica Kingsley Publishers; England, pp.152-161.

Davies S and Jasper M . (2004). A first-stage evaluation of a group programme for PND. *Community Practitioner*, 77(11), pp.426-433.

Fritz B and McGregor C . (2013). Postpartum depression screening in yemeni and punjabi immigrant women. *Nursing for Women's Health*, 17(3), pp.236-44.

Gordon T E and Cardone I A; Kim J J; Gordon S M; Silver R K;. (2006). Universal perinatal depression screening in an Academic Medical Center. *Obstetrics & Gynecology*, 107(2 Pt 1), pp.342-7.

Gutteridge K. (2001). The Tamworth Postnatal Depression Support Group. *MIDIRS Midwifery Digest*, 11(1), pp.S17-2.

Harmon Amy J. (2018). *Barriers to diagnosis of peripartum depression in Latino women.* . Harmon, Amy J.: William James College, Clinical Psychology, US.

Harris A L. (2017). Barriers to and Facilitators of Perinatal Depression Screening. *Nursing for Women's Health*, 20(6), pp.601-607.

Hayes B A. (2010). From 'postnatal depression' to 'perinatal anxiety and depression': key points of the National Perinatal Depression Plan for nurses and midwives in Australian primary health care settings. *Contemporary Nurse*, 35(1), pp.58-67.

Horowitz J A, Damato E and Solon L ; von Metzsch ; G ;. (1996). Identification of symptoms of postpartum depression: linking research to practice. *Journal of Perinatology*, 16(5), pp.360-5.

Howland L C, Jallo N and Connelly C D; Pickler R H;. (2017). Feasibility of a Relaxation Guided Imagery Intervention to Reduce Maternal Stress in the NICU. *JOGNN - Journal of Obstetric, Gynecologic and & Neonatal Nursing*, 46(4), pp.532-543.

Jallo N, Bourguignon C and Taylor A G; Ruiz J ; Goehler L ;. (2009). The biobehavioral effects of relaxation guided imagery on maternal stress. *Advances in Mind-Body Medicine*, 24(4), pp.12-22.

Kye Price, Sarah and Gray Lisa A; El-Khoury Dalia ;. (2014). Facilitating Mental Health Intervention in Home Visiting. *Zero to Three*, 34(5), pp.28-34.

Lehman Erica. (2012). *Centering pregnancy: A combined quantitative and qualitative appraisal of women's experiences of depression and anxiety during group prenatal care.* . Lehman, Erica: Wheaton Coll., US.

Lewis A, Ilot I and Lekka C ; Oluboyede Y ; (2011). Improving the quality of perinatal mental health: a health visitor-led protocol. *Community Practitioner*, 84(2), pp.27-31.

Liebert Heather. (2018). *Postpartum depression: Screening and treatment of low income women in a primary care setting.* . Liebert, Heather: The Wright Institute, US.

Lindsey Bruce, Daniel Béland and Angela Bowen ; (2012). MotherFirst: Developing a Maternal Mental Health Strategy in Saskatchewan. *Healthcare Policy*, 8(2), pp.46-55.

Marshall A, Guillen U and Mackley A ; Sturtz W ; (2019). Mindfulness Training among Parents with Preterm Neonates in the Neonatal Intensive Care Unit: A Pilot Study. *American Journal of Perinatology*, 36(14), pp.1514-1520.

McGregor Marla Louise. (2009). *A physician-based Cognitive Behavioral Intervention for depressed pregnant women in primary care: A pilot study.* . McGregor, Marla Louise: U Toronto, Canada.

Miller E S, Wisner K L; Gollan J and Hamade S ; Gossett D R; Grobman W A;. (2019). Screening and Treatment After Implementation of a Universal Perinatal Depression Screening Program. *Obstetrics & Gynecology*, 134(2), pp.303-309.

Mounts K O. (2009). Screening for maternal depression in the neonatal ICU. *Clinics in Perinatology*, 36(1), pp.137-52.

Owen L and Nevin H . (2008). Improving perinatal emotional well-being and mental health. *Nursing Times*, 104(8), pp.28-29.

Pearson Pamela, Klima Carrie and Snyder Marsha ; (2019). Reducing Barriers That Hinder Obstetric Providers From Addressing Perinatal Depression: A Provider Education Module. *Journal of Doctoral Nursing Practice*, 12(2), pp.212-224.

Poustie A and Drumm E . (1997). Co-ordinated care for women with postnatal depression. *Nursing Standard*, 11(16), pp.34-37.

Rastle Marsha. (2013). Screening for depression at first obstetrical visits at Naval Hospital Camp Pendleton in 2008: A comparison study of active duty members and military dependent wives with spouse's deployment status included. *Dissertation Abstracts International Section A: Humanities and Social Sciences*, 74(5-A(E)), pp.No Pagination Specified.

Robison Charlynn Phillis. (1999). *Effects of a stress management intervention on levels of stress and coping among pregnant and parenting adolescents: An investigation.(girls).* . Robison, Charlynn Phillis: Howard U., US.

Romain S, Jones A and Watts T ; (1995). Postnatal depression. Facilitating peer group support. *Health Visitor*, 68(4), pp.153-153.

Rowe Heather J, Calcagni Soledad Co; Galgut Sofia and Michelmores Janet ; Fisher Jane R;. (2014). Self-management of mild to moderate anxiety in women who have recently given

birth: Development and acceptability of a theoretically sound complex intervention. *International Journal of Mental Health Promotion*, 16(5), pp.308-319.

Rowe H J and Fisher J R. W;. (2015). Prevention of postnatal mental health problems in women: knowledge exchange in primary care in Victoria, Australia. *Health Promotion Journal of Australia*, 26(1), pp.64-69.

Stevens N R, Lillis T A; Wagner L and Tirone V ; Hobfoll S E;. (2019). A feasibility study of trauma-sensitive obstetric care for low-income, ethno-racial minority pregnant abuse survivors. *Journal of Psychosomatic Obstetrics & Gynecology*, 40(1), pp.66-74.

Tyldesley R and Phillips L . (2009). Beat the Blues -- a self-help group for parents experiencing postnatal depression. *A Life in the Day*, 13(2), pp.9-12.

Webster J, Linnane J and Dibley L ; Starrenburg S ; Roberts J ; Hinson J ;. (1997). The impact of screening for risk factors associated with postnatal depression at the first prenatal visit. *Journal of Quality in Clinical Practice*, 17(2), pp.65-71.

Wesley Y. (2006). Reduce stress: a stress reduction project for pregnant black women. *Journal of Cultural Diversity*, 13(4), pp.208-16.

Wilkinson J, Phillips S and Jackson J ; Walker K ;. (2003). "Mad for Fitness": an exercise group to combat a high incidence of postnatal depression. *Journal of Family Health Care*, 13(2), pp.44-8.

## **Not perinatal**

Bagnall Kara Marie. (2014). *Long-term follow-up of NetmumsHWD: A feasibility randomised controlled trial of telephone supported online behavioural activation for postnatal depression at 16 months post-randomisation*. . University of Exeter (United Kingdom).

Barlow Allison, Mullany Britta and Neault Nicole ; Goklish Novalene ; Billy Trudy ; Hastings Ranelda ; Lorenzo Sherilynn ; Kee Crystal ; Lake Kristin ; Redmond Cleve ; Carter Alice ; Walkup John T;. (2015). Paraprofessional-Delivered Home-Visiting Intervention for American Indian Teen Mothers and Children: 3-Year Outcomes From a Randomized Controlled Trial. *American Journal of Psychiatry*, 172(2), pp.154-162.

Beeber L S, Cooper C and Van Noy ; B E ; Schwartz T A; Blanchard H C; Canuso R ; Robb K ; Laudembacher C ; Emory S L;. (2007). Flying under the radar: engagement and retention of depressed low-income mothers in a mental health intervention. *Advances in Nursing Science*, 30(3), pp.221-34.

Cohen K, Leibovich L and Markin R ; Zilcha-Mano S ;. (2019). The process of change in psychotherapy with a pregnant patient following perinatal losses: An analysis of a case study. *Journal of clinical psychology*, 75(5), pp.874-885.

Cote-Arsenault D, Schwartz K and Krowchuk H ; McCoy T P;. (2014). Evidence-based intervention with women pregnant after perinatal loss. *MCN and American Journal of Maternal Child Nursing*, 39(3), pp.177-86; quiz 187-8.

Fletcher Richard, Campbell Linda and Sved Williams ; Anne ; Rawlinson Catherine ; Dye Jacob ; Baldwin Andrea ; May Chris ; StGeorge Jennifer ;. (2018). Sms4 perinatal parents:

Designing parenting support via text messages for mothers with severe mental illness (smi) and their partners. *Advances in Mental Health*, , pp.No Pagination Specified.

Volpe Ellen M, Quinn Camille R; Resch Kathryn and Douglas Valerie ; Cerulli Catherine ;. (2017). Assessing the feasibility and acceptability of Narrative Exposure Therapy to address IPV-related mental health in parenting and pregnant adolescents. *Journal of Family Violence*, 32(4), pp.439-452.

### **Not mental health**

Golfenshtein N, Deatrck J A and Lisanti A J; Medoff-Cooper B ;. (2017). Coping with the Stress in the Cardiac Intensive Care Unit: Can Mindfulness Be the Answer?. *Journal of Pediatric Nursing*, 37, pp.117-126.

Keller C, Vega-Lopez S and Ainsworth B ; Nagle-Williams A ; Records K ; Permana P ; Coonrod D ;. (2014). Social marketing: approach to cultural and contextual relevance in a community-based physical activity intervention. *Health Promotion International*, 29(1), pp.130-40.

Mason L, Roe B and Wong H ; Davies J ; Bamber J ;. (2010). The role of antenatal pelvic floor muscle exercises in prevention of postpartum stress incontinence: a randomised controlled trial. *Journal of Clinical Nursing*, 19(19-20), pp.2777-86.

Melnyk B M and Oswalt K L; Sidora-Arcoleo K. (2014). Validation and psychometric properties of the neonatal intensive care unit parental beliefs scale. *Nursing Research*, 63(2), pp.105-15.

Ngai F W and Wong P W; Chung K F; Leung K Y;. (2016). The effect of telephone-based cognitive-behavioural therapy on parenting stress: A randomised controlled trial. *Journal of Psychosomatic Research*, 86, pp.34-8.

Ngai F W and Wong P W; Chung K F; Leung K Y;. (2017). The effect of a telephone-based cognitive behavioral therapy on quality of life: a randomized controlled trial. *Archives of Women's Mental Health*, 20(3), pp.421-426.

Nolan A and Lawrence C . (2009). A pilot study of a nursing intervention protocol to minimize maternal-infant separation after Cesarean birth. *JOGNN - Journal of Obstetric, Gynecologic and Neonatal Nursing*, 38(4), pp.430-42.

O'Brien K, Bracht M and Macdonell K ; McBride T ; Robson K ; O'Leary L ; Christie K ; Galarza M ; Dicky T ; Levin A ; Lee S K;. (2013). A pilot cohort analytic study of Family Integrated Care in a Canadian neonatal intensive care unit. *BMC Pregnancy & Childbirth*, 13 Suppl 1, pp.S12.

Ong S L, Abdullah K L; Danaee M and Soh K L; Soh K G; Lee D S. K; Hussin E O. D;. (2019). The effectiveness of a structured nursing intervention program on maternal stress and ability among mothers of premature infants in a neonatal intensive care unit. *Journal of Clinical Nursing*, 28(3-4), pp.641-649.

Osman H, Saliba M and Chaaya M ; Naasan G ;. (2014). Interventions to reduce postpartum stress in first-time mothers: a randomized-controlled trial. *BMC Women's Health*, 14, pp.125.



Petteys A R and Adoumie D. (2018). Mindfulness-Based Neurodevelopmental Care: Impact on NICU Parent Stress and Infant Length of Stay; A Randomized Controlled Pilot Study. *Advances in Neonatal Care*, 18(2), pp.E12-E22.

Pouyan F, Kamrani M A and Rahimzadeh M ; Jamshidimanesh M ; Esmaelzadeh-Saeieh S ;. (2019). Effect of interpersonal psychotherapy oriented child birth education on stress and role adaptation in mothers with premature birth: A randomized clinical trial. *Iranian Journal of Psychiatry and Behavioral Sciences*, 13(1), pp..

Pulliaainen H, Niela-Vilen H and Ekholm E ; Ahlqvist-Bjorkroth S ;. (2019). Experiences of interactive ultrasound examination among women at risk of preterm birth: a qualitative study. *BMC Pregnancy & Childbirth*, 19(1), pp.338.

Reza N, Ali S M and Saeed K ; Abul-Qasim A ; Reza T H;. (2007). The impact of music on postoperative pain and anxiety following cesarean section. *Middle East Journal of Anesthesiology*, 19(3), pp.573-86.

Samra H A, Dutcher J and McGrath J M; Foster M ; Klein L ; Djira G ; Hansen J ; Wallenburg D ;. (2015). Effect of Skin-to-Skin Holding on Stress in Mothers of Late-Preterm Infants: A Randomized Controlled Trial. *Advances in Neonatal Care*, 15(5), pp.354-64.

Simavli S, Gumus I and Kaygusuz I ; Yildirim M ; Usluogullari B ; Kafali H ;. (2014). Effect of music on labor pain relief, anxiety level and postpartum analgesic requirement: A randomized controlled clinical trial. *Gynecologic and Obstetric Investigation*, 78(4), pp.244-250.

Snyder Kailey and Dinkel Danae . (2019). Mental and physical support (MAPS) for moms: preliminary findings from a prenatal health support program. *Integrative Medicine Research*, 8(1), pp.8-14.

Urizar G G, Jr and Yim I S; Rodriguez A ; Schetter C D;. (2019). The SMART Moms Program: A Randomized Trial of the Impact of Stress Management on Perceived Stress and Cortisol in Low-Income Pregnant Women. *Psychoneuroendocrinology*, 104, pp.174-184.

Urizar G G, Jr and Caliboso M ; Gearhart C ; Yim I S; Dunkel Schetter ; C ;. (2019). Process Evaluation of a Stress Management Program for Low-Income Pregnant Women: The SMART Moms/Mamas LISTAS Project. *Health Education & Behavior*, 46(6), pp.930-941.

Villamizar-Carvajal B, Vargas-Porras C and Garcia-Corzo J R;. (2018). Decrease of premature mothers' stress levels in the intensive care unit. *Enfermeria Intensiva*, 29(3), pp.113-120.

### **Substance use**

Kern J K and West E Y; Grannemann B D; Greer T L; Snell L M; Cline L L; VanBeveren T T; Heartwell S F; Kleiber B A; Trivedi M H;. (2004). Reductions in stress and depressive symptoms in mothers of substance-exposed infants, participating in a psychosocial program. *Maternal & Child Health Journal*, 8(3), pp.127-36.

### **No assessment/care/intervention**

- Abrams L S, Dornig K and Curran L ;. (2009). Barriers to service use for postpartum depression symptoms among low-income ethnic minority mothers in the United States. *Qualitative Health Research*, 19(4), pp.535-51.
- Abrams D, Nguyen L T and Murphy J ; Lee Y A; Tran N K; Wiljer D ;. (2016). Perceptions and experiences of perinatal mental disorders in rural, predominantly ethnic minority communities in northern Vietnam. *International Journal of Mental Health Systems*, 10, pp.10.
- Aftyka A, Rybojad B and Rosa W ; Wrobel A ; Karakula-Juchnowicz H ;. (2017). Risk factors for the development of post-traumatic stress disorder and coping strategies in mothers and fathers following infant hospitalisation in the neonatal intensive care unit. *Journal of Clinical Nursing*, 26(23-24), pp.4436-4445.
- Alfayumi-Zeadna S, Froimovici M and Azbarga Z ; Grotto I ; Daoud N ;. (2019). Barriers to postpartum depression treatment among Indigenous Bedouin women in Israel: A focus group study. *Health & Social Care in the Community*, 27(3), pp.757-766.
- Alhasanat Dalia, Fry - McComish and Judith ; Yarandi Hossein N;. (2017). Risk For Postpartum Depression Among Immigrant Arabic Women in the United States: A Feasibility Study. *Journal of Midwifery & Women's Health*, 62(4), pp.470-476.
- Allbaugh L J and Marcus S M; Ford E C; Flynn H A;. (2015). Development of a screening and recruitment registry to facilitate perinatal depression research in obstetrics settings in the USA. *International Journal of Gynaecology & Obstetrics*, 128(3), pp.260-3.
- Armstrong S J and Small R E;. (2010). The paradox of screening: rural women's views on screening for postnatal depression. *BMC Public Health*, 10, pp.744.
- Austin M P, Colton J and Priest S ; Reilly N ; Hadzi-Pavlovic D ;. (2013). The antenatal risk questionnaire (ANRQ): acceptability and use for psychosocial risk assessment in the maternity setting. *Women & Birth: Journal of the Australian College of Midwives*, 26(1), pp.17-25.
- Battle C L and Salisbury A L; Schofield C A; Ortiz-Hernandez S. (2013). Perinatal antidepressant use: Understanding women's preferences and concerns. *Journal of Psychiatric Practice*, 19(6), pp.443-453.
- Bayrampour H, McNeil D A and Benzies K ; Salmon C ; Gelb K ; Tough S ;. (2017). A qualitative inquiry on pregnant women's preferences for mental health screening. *BMC Pregnancy & Childbirth*, 17(1), pp.339.
- Bentley S M and Melville J L; Berry B D; Katon W J;. (2007). Implementing a clinical and research registry in obstetrics: overcoming the barriers. *General Hospital Psychiatry*, 29(3), pp.192-8.
- Bray N, Grasby K L and Lind P A; Painter J N; Colodro-Conde L ; Medland S E;. (2019). The psychosocial impact of nausea and vomiting during pregnancy as a predictor of postpartum depression. *Journal of Health Psychology*, , pp.1359105319859048.
- Byatt N, Cox L and Moore Simas ; T A ; Kini N ; Biebel K ; Sankaran P ; Swartz H A; Weinreb L ;. (2018). How obstetric settings can help address gaps in psychiatric care for pregnant and postpartum women with bipolar disorder. *Archives of Women's Mental Health*, 21(5), pp.543-551.

Chabrol H, Teissedre F and Armitage J ; Danel M ; Walburg V ;. (2004). Acceptability of psychotherapy and antidepressants for postnatal depression among newly delivered mothers. *Journal of Reproductive and Infant Psychology*, 22(1), pp.5-12.

Da Costa , D and Zelkowitz P ; Nguyen T V; Deville-Stoetzel J B;. (2018). Mental health help-seeking patterns and perceived barriers for care among nulliparous pregnant women. *Archives of Women's Mental Health*, 21(6), pp.757-764.

Dietsch E, Martin T and Shackleton P ; Davies C ; McLeod M ; Alston M ;. (2011). Australian Aboriginal kinship: a means to enhance maternal well-being. *Women & Birth: Journal of the Australian College of Midwives*, 24(2), pp.58-64.

El-Den S, O'Reilly C L and Chen T F;. (2019). Development and Psychometric Evaluation of a Questionnaire to Measure Attitudes Toward Perinatal Depression and Acceptability of Screening: The PND Attitudes and Screening Acceptability Questionnaire (PASAQ). *Evaluation & the Health Professions*, 42(4), pp.498-522.

Elkhodr S, Saba M and O'Reilly C ; Saini B ;. (2018). The role of community pharmacists in the identification and ongoing management of women at risk for perinatal depression: A qualitative study. *International Journal of Social Psychiatry*, 64(1), pp.37-48.

Evans K, Morrell C J and Spiby H ;. (2017). Women's views on anxiety in pregnancy and the use of anxiety instruments: a qualitative study. *Journal of Reproductive & Infant Psychology*, 35(1), pp.77-90.

Fonseca A, Gorayeb R and Canavarro M C;. (2015). Women's help-seeking behaviours for depressive symptoms during the perinatal period: Socio-demographic and clinical correlates and perceived barriers to seeking professional help. *Midwifery*, 31(12), pp.1177-85.

Foulkes Michelle. (2011). Enablers and Barriers to Seeking Help for a Postpartum Mood Disorder. *JOGNN: Journal of Obstetric and Gynecologic & Neonatal Nursing*, 40(4), pp.450-457.

Gemmill A W, Leigh B and Ericksen J ; Milgrom J ;. (2006). A survey of the clinical acceptability of screening for postnatal depression in depressed and non-depressed women. *BMC Public Health*, 6, pp.211.

Godderis R, Adair C E and Brager N ;. (2009). Applying new techniques to an old ally: a qualitative validation study of the Edinburgh Postnatal Depression Scale. *Women & Birth: Journal of the Australian College of Midwives*, 22(1), pp.17-23.

Goodman J H. (2009). Women's attitudes, preferences, and perceived barriers to treatment for perinatal depression. *Birth*, 36(1), pp.60-9.

Goodman S H, Dimidjian S and Williams K G;. (2013). Pregnant African American women's attitudes toward perinatal depression prevention. *Cultural Diversity & Ethnic Minority Psychology*, 19(1), pp.50-7.

Gordon M, Henderson R and Holmes J H; Wolters M K; Bennett I M; Group Spirit ;. (2016). Participatory design of ehealth solutions for women from vulnerable populations with perinatal depression. *Journal of the American Medical Informatics Association*, 23(1), pp.105-9.

Goyal Deepika, Van Ta and Park ; McNiesh Susan ;. (2015). Postpartum Depression. *MCN: The American Journal of Maternal Child Nursing*, 40(4), pp.256-261.

Grissette B G, Spratling R and Aycock D M;. (2018). Barriers to Help-Seeking Behavior Among Women With Postpartum Depression. *JOGNN - Journal of Obstetric, Gynecologic and Neonatal Nursing*, 47(6), pp.812-819.

Guintivano J, Krohn H and Lewis C ; Byrne E M; Henders A K; Ploner A ; Kirk K ; Martin N G; Milgrom J ; Wray N R; Sullivan P F; Meltzer-Brody S ;. (2018). PPD ACT: an app-based genetic study of postpartum depression. *Translational Psychiatry*, 8(1), pp..

Henshaw E, Sabourin B and Warning M ;. (2013). Treatment-seeking behaviors and attitudes survey among women at risk for perinatal depression or anxiety. *JOGNN - Journal of Obstetric, Gynecologic and Neonatal Nursing*, 42(2), pp.168-77.

Holden G, Corter A L and Hatters-Friedman S ; Soosay I ;. (2019). Brief Report. A qualitative study of maternal mental health services in New Zealand: Perspectives of Maori and Pacific mothers and midwives. *Asia-Pacific Psychiatry*, (no pagination), pp..

Jarrett Patricia M. (2016). Pregnant women's experience of depression care. *Journal of Mental Health Training and Education & Practice*, 11(1), pp.33-47.

Jesse D E and Dolbier C L; Blanchard A. (2008). Barriers to seeking help and treatment suggestions for prenatal depressive symptoms: focus groups with rural low-income women. *Issues in Mental Health Nursing*, 29(1), pp.3-19.

Jomeen J, Glover L and Jones C ; Garg D ; Marshall C ;. (2013). Assessing women's perinatal psychological health: Exploring the experiences of health visitors. *Journal of Reproductive and Infant Psychology*, 31(5), pp.479-489.

Khajehpour M, Simbar M and Jannesari S ; Ramezani-Tehrani F ; Majd H A;. (2013). Health status of women with intended and unintended pregnancies. *Public Health*, 127(1), pp.58-64.

Kingston D, McDonald S and Tough S ; Austin M P; Hegadoren K ; Lasiuk G ;. (2014). Public views of acceptability of perinatal mental health screening and treatment preference: a population based survey. *BMC Pregnancy & Childbirth*, 14, pp.67.

Kingston D E, Biringer A and McDonald S W; Heaman M I; Lasiuk G C; Hegadoren K M; McDonald S D; Veldhuyzen van Zanten; S ; Sword W ; Kingston J J; Jarema K M; Vermeyden L ; Austin M P;. (2015). Preferences for Mental Health Screening Among Pregnant Women: A Cross-Sectional Study. *American Journal of Preventive Medicine*, 49(4), pp.e35-43.

Kinser P and Masho S . (2015). "I just start crying for no reason": the experience of stress and depression in pregnant, urban, African-American adolescents and their perception of yoga as a management strategy. *Womens Health Issues*, 25(2), pp.142-8.

Lara-Cinisomo S, Wisner K L and Burns R M; Chaves-Gnecco D ;. (2014). Perinatal depression treatment preferences among Latina mothers. *Qualitative Health Research*, 24(2), pp.232-41.

- Lasater Molly Elizabeth. (2019). Integrating mental health into community-based maternal health services in Selingue, Mali. *Dissertation Abstracts International: Section B: The Sciences and Engineering*, 80(8-B(E)), pp.No Pagination Specified.
- Leung S, Arthur D G and Martinson I ;. (2005). Stress in women with postpartum depression: a phenomenological study. *Journal of Advanced Nursing*, 51(4), pp.353-60.
- Logsdon M C, Hines-Martin V and Rakestraw V ;. (2009). Barriers to depression treatment in low-income, unmarried, adolescent mothers in a southern, urban area of the United States. *Issues in Mental Health Nursing*, 30(7), pp.451-5.
- Lowenhoff Catherine, Davison-Fischer Jan and Pike Nick ; Appleton Jane V;. (2019). Using the TIDieR checklist to describe health visitor support for mothers with mental health problems: Analysis of a cross-sectional survey. *Health & Social Care in the Community*, 27(5), pp.e824-e836.
- Massoudi P, Wickberg B and Hwang P ;. (2007). Screening for postnatal depression in Swedish child health care. *Acta Paediatrica*, 96(6), pp.897-901.
- McGookin A, Furber C and Smith D M;. (2017). Student midwives' awareness, knowledge, and experiences of antenatal anxiety within clinical practice. *Journal of Reproductive & Infant Psychology*, 35(4), pp.380-393.
- Mohamed H A, Spencer S L; Al Swasy and A H ; Swidan S E; Abouelenien M S;. (2014). A social and Biological Approach for Postpartum Depression in Egypt. *Woman - Psychosomatic Gynaecology and Obstetrics*, 1(C), pp.30-39.
- Moore D and Harrison V . (2018). Advice for Health Care Professionals and Users: An Evaluation of Websites for Perinatal Anxiety. *JMIR Mental Health*, 5(4), pp.e11464.
- Musyimi C W and Mutiso V N; Nyamai D N; Ebuenyi I D; Ndetei D M;. (2019). Integration of Traditional Birth Attendants into Mental Healthcare: A Multistakeholder Qualitative Study Exploration. *BioMed Research International*, 2019, pp.8195267.
- Muzik Maria, Kirk Rosalind and Alfafara Emily ; Jonika Jennifer ; Waddell Rachel ;. (2016). Teenage mothers of black and minority ethnic origin want access to a range of mental and physical health support: a participatory research approach. *Health Expectations*, 19(2), pp.403-415.
- Mwape L, McGuinness T M and Dixey R ; Johnson S E;. (2012). Socio-cultural factors surrounding mental distress during the perinatal period in Zambia: a qualitative investigation. *International Journal of Mental Health Systems*, 6(1), pp.12.
- Negron R, Martin A and Almog M ; Balbierz A ; Howell E A;. (2013). Social support during the postpartum period: mothers' views on needs, expectations, and mobilization of support. *Maternal & Child Health Journal*, 17(4), pp.616-23.
- Ng'oma M, Meltzer-Brody S and Chirwa E ; Stewart R C;. (2019). "Passing through difficult times": Perceptions of perinatal depression and treatment needs in Malawi - A qualitative study to inform the development of a culturally sensitive intervention. *PLoS ONE [Electronic Resource]*, 14(6), pp.e0217102.

- Nyberg K, Lindberg I and Ohrling K ;. (2010). Midwives' experience of encountering women with posttraumatic stress symptoms after childbirth. *Sexual & reproductive healthcare : official journal of the Swedish Association of Midwives*, 1(2), pp.55-60.
- Nygaard L, Rossen C B and Buus N ;. (2015). Balancing Risk: A Grounded Theory Study of Pregnant Women's Decisions to (Dis)Continue Antidepressant Therapy. *Issues in Mental Health Nursing*, 36(7), pp.485-92.
- O'Mahen H, Fedock G and Henshaw E ; Himle J A; Forman J ; Flynn H A;. (2012). Modifying CBT for Perinatal Depression: What Do Women Want?. A Qualitative Study. *Cognitive and Behavioral Practice*, 19(2), pp.359-371.
- O'Mahony J M, Donnelly T T; Raffin Bouchal and S ; Este D ;. (2012). Barriers and facilitators of social supports for immigrant and refugee women coping with postpartum depression. *Advances in Nursing Science*, 35(3), pp.E42-56.
- O'Mahony Joyce Maureen, Donnelly Tam Truong; Este Dave and Bouchal Shelley Raffin;. (2012). Using Critical Ethnography to Explore Issues among Immigrant and Refugee Women Seeking Help for Postpartum Depression. *Issues in Mental Health Nursing*, 33(11), pp.735-742.
- Oh S, Salas-Wright C P and Vaughn M G;. (2018). Trends in depression among low-income mothers in the United States, 2005-2015. *Journal of Affective Disorders*, 235, pp.72-75.
- Olin S S, McCord M and Stein R E. K; Kerker B D; Weiss D ; Hoagwood K E; Horwitz S M;. (2017). Beyond Screening: A Stepped Care Pathway for Managing Postpartum Depression in Pediatric Settings. *Journal of Women's Health*, 26(9), pp.966-975.
- Olson A L and Kemper K J; Kelleher K J; Hammond C S; Zuckerman B S; Dietrich A J;. (2002). Primary care pediatricians' roles and perceived responsibilities in the identification and management of maternal depression. *Pediatrics*, 110(6), pp.1169-76.
- Osok J, Kigamwa P and Huang K Y; Grote N ; Kumar M ;. (2018). Adversities and mental health needs of pregnant adolescents in Kenya: identifying interpersonal, practical, and cultural barriers to care. *BMC Women's Health*, 18(1), pp.96.
- Pawils S, Metzner F and Wendt C ; Raus S ; Shedden-Mora M ; Wlodarczyk O ; Harter M ;. (2016). Patients with Postpartum Depression in Gynaecological Practices in Germany - Results of a Representative Survey of Local Gynaecologists about Diagnosis and Management. *Geburtshilfe und Frauenheilkunde*, 76(8), pp.888-894.
- Pilkington P, Milne L and Cairns K ; Whelan T ;. (2016). Enhancing reciprocal partner support to prevent perinatal depression and anxiety: a Delphi consensus study. *BMC Psychiatry*, 16, pp.23.
- Place J M. S, Allen-Leigh B and Billings D L; Dues K M; de Castro ; F ;. (2017). Detection and care practices for postpartum depressive symptoms in public-sector obstetric units in Mexico: Qualitative results from a resource-constrained setting. *Birth*, 44(4), pp.390-396.
- Ransing R, Kukreti P and Deshpande S ; Godake S ; Neelam N ; Raghuveer P ; Mahadevaiah M ; Kataria D ; Patil S ; Puri M ; Padma K ;. (2019). Perinatal depression-knowledge gap among service providers and service utilizers in India. *Asian Journal of Psychiatry*, 47, pp.101822.

Raymond N C, Pratt R J; Godecker A and Harrison P A; Kim H ; Kuendig J ; O'Brien J M;. (2014). Addressing perinatal depression in a group of underserved urban women: a focus group study. *BMC Pregnancy & Childbirth*, 14, pp.336.

Recto P and Champion J D. (2018). "We Don't Want to be Judged": Perceptions about Professional Help and Attitudes Towards Help-Seeking among Pregnant and Postpartum Mexican-American Adolescents. *Journal of Pediatric Nursing*, 42, pp.111-117.

Reilly N, Harris S and Loxton D ; Chojenta C ; Forder P ; Milgrom J ; Austin M P;. (2013). Disparities in reported psychosocial assessment across public and private maternity settings: a national survey of women in Australia. *BMC Public Health*, 13, pp.632.

Rompala K S, Cirino N and Rosenberg K D; Fu R ; Lambert W E;. (2016). Prenatal Depression Screening by Certified Nurse-Midwives, Oregon. *Journal of Midwifery & Women's Health*, 61(5), pp.599-605.

Sanders L B. (2006). Attitudes, perceived ability, and knowledge about depression screening: a survey of certified nurse-midwives/certified midwives. *Journal of Midwifery & Women's Health*, 51(5), pp.340-346.

Segre L S, O'Hara M W; Arndt S and Beck C T;. (2010). Nursing care for postpartum depression, part 1: do nurses think they should offer both screening and counseling?. *MCN and American Journal of Maternal Child Nursing*, 35(4), pp.220-5.

Segre L S, O'Hara M W; Arndt S and Beck C T;. (2010). Screening and counseling for postpartum depression by nurses: the women's views. *MCN and American Journal of Maternal Child Nursing*, 35(5), pp.280-5.

Silverwood V, Nash A and Chew-Graham C A; Walsh-House J ; Sumathipala A ; Bartlam B ; Kingstone T ;. (2019). Healthcare professionals' perspectives on identifying and managing perinatal anxiety: a qualitative study. *British Journal of General Practice*, 69(688), pp.e768-e776.

Skocir A P and Hundley V. (2006). Are Slovenian midwives and nurses ready to take on a greater role in caring for women with postnatal depression?. *Midwifery*, 22(1), pp.40-55.

Skoog M, Hallstrom I and Berggren V ;. (2017). 'There's something in their eyes' - Child Health Services nurses' experiences of identifying signs of postpartum depression in non-Swedish-speaking immigrant mothers. *Scandinavian Journal of Caring Sciences*, 31(4), pp.739-747.

Sleath B, West S and Tudor G ; Perreira K ; King V ; Morrissey J ;. (2005). Ethnicity and depression treatment preferences of pregnant women. *Journal of Psychosomatic Obstetrics & Gynecology*, 26(2), pp.135-40.

Surjaningrum E R, Minas H and Jorm A F; Kakuma R ;. (2018). The feasibility of a role for community health workers in integrated mental health care for perinatal depression: a qualitative study from Surabaya, Indonesia. *International Journal of Mental Health Systems*, 12, pp.27.

Tabb K M, Choi S and Pinerros-Leano M ; Meline B ; McDonald H G; Kester R ; Huang H ;. (2015). Perinatal depression screening in a Women, Infants, and Children (WIC) program:

perception of feasibility and acceptability among a multidisciplinary staff. *General Hospital Psychiatry*, 37(4), pp.305-9.

Teng L, Robertson Blackmore and E ; Stewart D E;. (2007). Healthcare worker's perceptions of barriers to care by immigrant women with postpartum depression: an exploratory qualitative study. *Archives of Women's Mental Health*, 10(3), pp.93-101.

Thomas N, Komiti A and Judd F ;. (2014). Pilot early intervention antenatal group program for pregnant women with anxiety and depression. *Archives of Women's Mental Health*, 17(6), pp.503-9.

Thomas L J and Scharp K M; Paxman C G;. (2014). Stories of postpartum depression: exploring health constructs and help-seeking in mothers' talk. *Women & Health*, 54(4), pp.373-87.

Tol W A, Ebrecht B and Aiyo R ; Murray S M; Nguyen A J; Kohrt B A; Ndyanabangi S ; Alderman S ; Musisi S ; Nakku J ;. (2018). Maternal mental health priorities, help-seeking behaviors, and resources in post-conflict settings: a qualitative study in eastern Uganda. *BMC Psychiatry*, 18(1), pp.39.

Vanderkruik Rachel and Dimidjian Sona . (2019). Perspectives on task-shifting depression care to peers for depressed Latina mothers. *Journal of Latinx Psychology*, 7(1), pp.22-38.

Vigod S, Sultana A and Fung K ; Hussain-Shamsy N ; Dennis C L;. (2016). A Population-Based Study of Postpartum Mental Health Service Use by Immigrant Women in Ontario, Canada. *Canadian Journal of Psychiatry - Revue Canadienne de Psychiatrie*, 61(11), pp.705-713.

Walker L O and Im E O; Tyler D O;. (2013). Maternal health needs and interest in screening for depression and health behaviors during pediatric visits. *Journal of Pediatric Health Care*, 27(4), pp.267-77.

Wenze S J and Battle C L;. (2018). Perinatal Mental Health Treatment Needs, Preferences, and Barriers in Parents of Multiples. *Journal of Psychiatric Practice*, 24(3), pp.158-168.

Zittel-Palamara Kimberley, Rockmaker Julie R and Schwabel Kara M; Weinstein Wendy L; Thompson Sanna J;. (2008). Desired assistance versus care received for postpartum depression: access to care differences by race. *Archives of Women's Mental Health*, 11(2), pp.81-92.

## **Outcome**

Abdollahpour S, Khosravi A and Bolbolhaghghi N ;. (2016). The effect of the magical hour on post-traumatic stress disorder (PTSD) in traumatic childbirth: a clinical trial. *Journal of Reproductive and Infant Psychology*, 34(4), pp.403-412.

Abdollahpour S, Khosravi A and Motaghi Z ; Keramat A ; Mousavi S A;. (2019). Effect of Brief Cognitive Behavioral Counseling and Debriefing on the Prevention of Post-traumatic Stress Disorder in Traumatic Birth: A Randomized Clinical Trial. *Community Mental Health Journal*, 55(7), pp.1173-1178.



- Abdollahpour S, Bolbolhaghghi N and Khosravi A ;. (2019). Effect of the Sacred Hour on Postnatal Depression in Traumatic Childbirth: a Randomized Controlled Trial. *Journal of Caring Sciences*, 8(2), pp.69-74.
- Abiodun O A. (2006). Postnatal depression in primary care populations in Nigeria. *General Hospital Psychiatry*, 28(2), pp.133-6.
- Abrahams Z, Schneider M and Field S ; Honikman S ;. (2019). Validation of a brief mental health screening tool for pregnant women in a low socio-economic setting. *BMC psychology*, 7(1), pp.77.
- Adams E and Tongs J . (2011). Starting a perinatal and infant mental health service at Winnunga Nimmityjah. *Australasian Psychiatry*, 19 Suppl 1, pp.S20-2.
- Ahlqvist-Bjorkroth S, Axelin A and Korja R ; Lehtonen L ;. (2019). An educational intervention for NICU staff decreased maternal postpartum depression. *Pediatric Research*, 85(7), pp.982-986.
- Ahn Y M and Kim M R;. (2004). The effects of a home-visiting discharge education on maternal self-esteem, maternal attachment, postpartum depression and family function in the mothers of NICU infants. *Daehan Ganho Haghoeji*, 34(8), pp.1468-76.
- Aite L, Trucchi A and Nahom A ; Spina V ; Bilancioni E ; Bagolan P ;. (2002). Multidisciplinary management of fetal surgical anomalies: the impact on maternal anxiety. *European Journal of Pediatric Surgery*, 12(2), pp.90-4.
- Aite L, Trucchi A and Nahom A ; Zaccara A ; Casaccia G ; Bagolan P ;. (2003). A challenging intervention with maternal anxiety: Babies requiring surgical correction of a congenital anomaly after missed prenatal diagnosis. *Infant Mental Health Journal*, 24(6), pp.571-579.
- Akbarzadeh M, Rad S K and Moattari M ; Zare N ;. (2018). Investigation of breastfeeding training based on BASNEF model on the intensity of postpartum blues. *Eastern Mediterranean Health Journal*, 23(12), pp.830-835.
- Alder E and Truman J . (2002). Counselling for postnatal depression in the voluntary sector. *Psychology & Psychotherapy: Theory and Research & Practice*, 75(Pt 2), pp.207-20.
- Alder E M, Reid M and Sharp L J; Cantwell R ; Robertson K ; Kearney E ;. (2008). Policy and practice in the management of postnatal depression in Scotland. *Archives of Women's Mental Health*, 11(3), pp.213-9.
- Alexander P. (2013). Innovation:CBT-based support groups for postnatal depression. *Nursing Times*, 109(41), pp.12-4.
- Ali N S and Ali B S; Azam I S;. (2009). Post partum anxiety and depression in peri-urban communities of Karachi, Pakistan: a quasi-experimental study. *BMC Public Health*, 9, pp.384.
- Ammerman R T, Peugh J L; Putnam F W; Van Ginkel and J B ;. (2012). Predictors of treatment response in depressed mothers receiving in-home cognitive-behavioral therapy and concurrent home visiting. *Behavior Modification*, 36(4), pp.462-81.

Ammerman R T, Putnam F W; Altaye M and Teeters A R; Stevens J ; Van Ginkel ; J B ;. (2013). Treatment of depressed mothers in home visiting: impact on psychological distress and social functioning. *Child Abuse & Neglect*, 37(8), pp.544-54.

Ammerman R T, Putnam F W; Altaye M and Stevens J ; Teeters A R; Van Ginkel ; J B ;. (2013). A clinical trial of in-home CBT for depressed mothers in home visitation. *Behavior Therapy*, 44(3), pp.359-72.

Ammerman R T, Altaye M and Putnam F W; Teeters A R; Zou Y ; Van Ginkel ; J B ;. (2015). Depression improvement and parenting in low-income mothers in home visiting. *Archives of Women's Mental Health*, 18(3), pp.555-63.

Trevillion K, Ryan E G and Pickles A ; Heslin M ; Byford S ; Nath S ; Bick D ; Milgrom J ; Mycroft R ; Domoney J ; Pariante C ; Hunter M S; Howard L M;. ( ). An exploratory parallel-group randomised controlled trial of antenatal Guided Self-Help (plus usual care) versus usual care alone for pregnant women with depression: DAWN trial. *Journal of Affective Disorders*, 261, pp.187-197.

Anagnostaki L, Kollia I and Layiou-Lignos E ;. (2019). Implementation of a brief early intervention in times of socio-economic crisis: effects on parental stress. *Journal of Child Psychotherapy*, 45(1), pp.55-70.

Ancill R, Hilton S and Carr T ; Tooley M ; McKenzie A ;. (1986). Screening for antenatal and postnatal depressive symptoms in general practice using a microcomputer-delivered questionnaire. *Journal of the Royal College of General Practitioners*, 36(287), pp.276-9.

Anderson Carla M and Brunton Robyn J; Dryer Rachel. (2018). Pregnancy-related anxiety: Re-examining its distinctiveness+. *Australian Psychologist*, , pp.No Pagination Specified.

Anderson C A and Connolly J P;. (2018). Predicting posttraumatic stress and depression symptoms among adolescents in the extended postpartum period. *Heliyon*, 4(11), pp.e00965.

Appleby L, Hirst E and Marshall S ; Keeling F ; Brind J ; Butterworth T ; Lole J ;. (2003). The treatment of postnatal depression by health visitors: impact of brief training on skills and clinical practice. *Journal of Affective Disorders*, 77(3), pp.261-6.

Armstrong K L and Fraser J A; Dadds M R; Morris J. (1999). A randomized, controlled trial of nurse home visiting to vulnerable families with newborns. *Journal of Paediatrics & Child Health*, 35(3), pp.237-44.

Armstrong K and Edwards H . (2004). The effectiveness of a pram-walking exercise programme in reducing depressive symptomatology for postnatal women. *International Journal of Nursing Practice*, 10(4), pp.177-94.

Armstrong S and Small R . (2007). Screening for postnatal depression: not a simple task. *Australian & New Zealand Journal of Public Health*, 31(1), pp.57-61.

Arroyo Carl and Fowler Neil . (2013). Before and after: A mother and infant painting group. *International Journal of Art Therapy*, 18(3), pp.98-112.

Ashby B, Ranadive N and Alaniz V ; St John-Larkin ; C ; Scott S ;. (2016). Implications of Comprehensive Mental Health Services Embedded in an Adolescent Obstetric Medical Home. *Maternal & Child Health Journal*, 20(6), pp.1258-65.

Ashby B D and Ehmer A C; Scott S M;. (2019). Trauma-informed care in a patient-centered medical home for adolescent mothers and their children. *Psychological Services*, 16(1), pp.67-74.

Ashford M T, Olander E K; Rowe H and Fisher J R; Ayers S ;. (2018). Feasibility and Acceptability of a Web-Based Treatment with Telephone Support for Postpartum Women With Anxiety: Randomized Controlled Trial. *JMIR Mental Health*, 5(2), pp.e19.

Atif N, Krishna R N and Sikander S ; Lazarus A ; Nisar A ; Ahmad I ; Raman R ; Fuhr D C; Patel V ; Rahman A ;. (2017). Mother-to-mother therapy in India and Pakistan: adaptation and feasibility evaluation of the peer-delivered Thinking Healthy Programme. *BMC Psychiatry*, 17(1), pp.79.

Atif N, Nisar A and Bibi A ; Khan S ; Zulfiqar S ; Ahmad I ; Sikander S ; Rahman A ;. (2019). Scaling-up psychological interventions in resource-poor settings: training and supervising peer volunteers to deliver the 'Thinking Healthy Programme' for perinatal depression in rural Pakistan. *Global Mental Health*, 6, pp.e4.

Atif N, Bibi A and Nisar A ; Zulfiqar S ; Ahmed I ; LeMasters K ; Hagaman A ; Sikander S ; Maselko J ; Rahman A ;. (2019). Delivering maternal mental health through peer volunteers: a 5-year report. *International Journal of Mental Health Systems*, 13, pp.62.

Avalos L A, Raine-Bennett T and Chen H ; Adams A S; Flanagan T ;. (2016). Improved Perinatal Depression Screening, Treatment, and Outcomes With a Universal Obstetric Program. *Obstetrics & Gynecology*, 127(5), pp.917-25.

Ayers S, Claypool J and Eagle A ;. (2006). Birth trauma. What happens after a difficult birth? Postnatal debriefing services. *British Journal of Midwifery*, 14(3), pp.157-161.

Ayers S, Fitzgerald G and Thompson S ;. (2015). Brief Online Self-help Exercises for Postnatal Women to Improve Mood: A Pilot Study. *Maternal & Child Health Journal*, 19(11), pp.2375-83.

Babu G R, Murthy G V. S; Singh N and Nath A ; Rathnaiah M ; Saldanha N ; Deepa R ; Kinra S ;. (2018). Sociodemographic and Medical Risk Factors Associated With Antepartum Depression. *Frontiers in Public Health*, 6, pp.127.

Bakare M O and Bello-Mojeed M A; Munir K M; Duduyemi O O; Orowigho A O; Odetunde O I; Taiwo O G; Olofinlade J A; Omotoso O N; Famurewa O H; Omolabi O O; Jejeloye A O;. (2017). Improving access to interventions among mothers screened positive for post-partum depression (PPD) at National Programme on Immunization (NPI) clinics in south-western and south-eastern Nigeria - A service development report. *Matters*, , pp.

Baker L, Cross S and Greaver L ; Wei G ; Lewis R ; Healthy Start ; Corps ;. (2005). Prevalence of postpartum depression in a native American population. *Maternal & Child Health Journal*, 9(1), pp.21-5.

Baker C D, Kamke H and O'Hara M W; Stuart S ;. (2009). Web-based training for implementing evidence-based management of postpartum depression. *Journal of the American Board of Family Medicine: JABFM*, 22(5), pp.588-9.

Baker-Ericzen Mary J, Mueggenborg Mary Garnand; Hartigan Phyllis and Howard Nicole ; Wilke Terry ;. (2008). Partnership for women's health: A new-age collaborative program for addressing maternal depression in the postpartum period. *Families, Systems and & Health*, 26(1), pp.30-43.

Baker-Ericzen M J, Connelly C D; Hazen A L; Duenas C and Landsverk J A; Horwitz S M;. (2012). A collaborative care telemedicine intervention to overcome treatment barriers for Latina women with depression during the perinatal period. *Families, Systems and & Health*, 30(3), pp.224-40.

Balbino F S and Balieiro M M; Mandetta M A;. (2016). Measurement of Family-centered care perception and parental stress in a neonatal unit. *Revista Latino-Americana de Enfermagem*, 24, pp.e2753.

Baldisserotto M L, Miranda Theme and M ; Gomez L Y; Dos Reis ; T B Q;. (2019). Barriers to Seeking and Accepting Treatment for Perinatal Depression: A Qualitative Study in Rio de Janeiro, Brazil. *Community Mental Health Journal*, 10, pp.10.

Bansal S C and Ganjiwale J D; Nimbalkar S M; Kharod N M;. (2018). Screening for postpartum depression in a neonatal intensive care unit and postnatal ward with its impact on newborn care practices: A hospital based survey. *Journal of Clinical and Diagnostic Research*, 12(8), pp.VC01-VC06.

Barnes J, Senior R and MacPherson K ;. (2009). The utility of volunteer home-visiting support to prevent maternal depression in the first year of life. *Child: Care and Health & Development*, 35(6), pp.807-16.

Barnet B, Liu J and DeVoe M ; Alperovitz-Bichell K ; Duggan A K;. (2007). Home visiting for adolescent mothers: effects on parenting, maternal life course, and primary care linkage. *Annals of Family Medicine*, 5(3), pp.224-32.

Barnett B, Matthey S and Gyaneshwar R ;. (1999). Screening for postnatal depression in women of non-English speaking background. *Archives of Women's Mental Health*, 2(2), pp.67-74.

Barrera A Z and Wickham R E; Munoz R F;. (2015). Online prevention of postpartum depression for Spanish- and English-speaking pregnant women: A pilot randomized controlled trial. *Internet Interventions*, 2(3), pp.257-265.

Barrera A Z, Aguilera A and Inlow N ; Servin J ;. (2019). A preliminary study on the acceptability of a brief SMS program for perinatal women. *Health Informatics Journal*, , pp.1460458219866560.

Barry Leasha M and Singer George H;. (2001). Reducing maternal psychological distress after the NICU experience through journal writing. *Journal of Early Intervention*, 24(4), pp.287-297.

- Battle C L and Howard M M;. (2014). A mother-baby psychiatric day hospital: History, rationale, and why perinatal mental health is important for obstetric medicine. *Obstetric Medicine*, 7(2), pp.66-70.
- Battle C L and Uebelacker L A; Magee S R; Sutton K A; Miller I W;. (2015). Potential for prenatal yoga to serve as an intervention to treat depression during pregnancy. *Womens Health Issues*, 25(2), pp.134-41.
- Bauer C L, Victorson D and Rosenbloom S ; Barocas J ; Silver R K;. (2010). Alleviating distress during antepartum hospitalization: a randomized controlled trial of music and recreation therapy. *Journal of Women's Health*, 19(3), pp.523-31.
- Baumel A, Tinkelman A and Mathur N ; Kane J M;. (2018). Digital Peer-Support Platform (7Cups) as an Adjunct Treatment for Women With Postpartum Depression: Feasibility, Acceptability, and Preliminary Efficacy Study. *JMIR MHealth and UHealth*, 6(2), pp.e38.
- Beattie J, Hall H and Biro M A; East C ; Lau R ;. (2017). Effects of mindfulness on maternal stress, depressive symptoms and awareness of present moment experience: A pilot randomised trial. *Midwifery*, 50, pp.174-183.
- Beckjord E B, Keyser D J; Schultz D and Lovejoy S L; Firth R ; Pincus H A;. (2011). Potential benefits of health information technology for integrating physical and behavioral health care: perinatal depression as a case-in-point. *Translational Behavioral Medicine*, 1(1), pp.89-92.
- Beddoe Amy E. (2008). Mindfulness-based yoga during pregnancy: A pilot study examining relationships between stress, anxiety, sleep, and pain. *Dissertation Abstracts International: Section B: The Sciences and Engineering*, 68(7-B), pp.4384.
- Beddoe A E, Paul Yang and C P ; Kennedy H P; Weiss S J; Lee K A;. (2009). The effects of mindfulness-based yoga during pregnancy on maternal psychological and physical distress. *JOGNN - Journal of Obstetric, Gynecologic and Neonatal Nursing*, 38(3), pp.310-9.
- Beeber L S, Holditch-Davis D and Perreira K ; Schwartz T A; Lewis V ; Blanchard H ; Canuso R ; Goldman B D;. (2010). Short-term in-home intervention reduces depressive symptoms in Early Head Start Latina mothers of infants and toddlers. *Research in Nursing & Health*, 33(1), pp.60-76.
- Beevi Z, Low W Y and Hassan J ;. (2016). Impact of Hypnosis Intervention in Alleviating Psychological and Physical Symptoms During Pregnancy. *American Journal of Clinical Hypnosis*, 58(4), pp.368-82.
- Bell Linda, Feeley Nancy and Hayton Barbara ; Zekowitz Phyllis ; Tait Madeleine ; Desindes Sophie ;. (2016). Barriers and Facilitators to the Use of Mental Health Services by Women With Elevated Symptoms of Depression and Their Partners. *Issues in Mental Health Nursing*, 37(9), pp.651-659.
- Benediktsson I, McDonald S W and Vekved M ; McNeil D A; Dolan S M; Tough S C;. (2013). Comparing CenteringPregnancy to standard prenatal care plus prenatal education. *BMC Pregnancy & Childbirth*, 13 Suppl 1, pp.S5.
- Berard A, Sheehy O and Zhao J P; Chambers C ; Roth M ; Bozzo P ; Johnson D ; Kao K ; Lavigne S ; Wolfe L ; Quinn D ; Dieter K ; MotherToBaby Collaborative Research;

Committee ;. (2019). Impact of antidepressant use, discontinuation, and dosage modification on maternal depression during pregnancy. *European Neuropsychopharmacology*, 29(7), pp.803-812.

Bershadsky S, Trumppheller L and Kimble H B; Pipaloff D ; Yim I S;. (2014). The effect of prenatal Hatha yoga on affect, cortisol and depressive symptoms. *Complementary Therapies in Clinical Practice*, 20(2), pp.106-13.

Bhat A, Reed S and Mao J ; Vredevoogd M ; Russo J ; Unger J ; Rowles R ; Unutzer J ;. (2018). Delivering perinatal depression care in a rural obstetric setting: a mixed methods study of feasibility, acceptability and effectiveness. *Journal of Psychosomatic Obstetrics & Gynecology*, 39(4), pp.273-280.

Bhat A, Mao J and Unutzer J ; Reed S ; Unger J ;. (2018). Text messaging to support a perinatal collaborative care model for depression: A multi-methods inquiry. *General Hospital Psychiatry*, 52, pp.14-20.

Bhusal B R, Bhandari N and Chapagai M ; Gavidia T ;. (2016). Validating the Edinburgh Postnatal Depression Scale as a screening tool for postpartum depression in Kathmandu, Nepal. *International Journal of Mental Health Systems*, 10, pp.71.

Biggs L J, Shafiei T and Forster D A; Small R ; McLachlan H L;. (2015). Exploring the views and experiences of callers to the PANDA Post and Antenatal Depression Association Australian National Perinatal Depression Helpline: a cross-sectional survey. *BMC Pregnancy & Childbirth*, 15, pp.209.

Biggs Laura J, McLachlan Helen L; Shafiei Touran and Small Rhonda ; Forster Della A;. (2019). Peer supporters' experiences on an Australian perinatal mental health helpline. *Health Promotion International*, 34(3), pp.479-489.

Biggs L J, McLachlan H L; Shafiei T and Liamputtong P ; Forster D A;. (2019). 'I need help': Reasons new and re-engaging callers contact the PANDA-Perinatal Anxiety and Depression Australia National Helpline. *Health & Social Care in the Community*, 27(3), pp.717-728.

Bina R, Barak A and Posmontier B ; Glasser S ; Cinamon T ;. (2019). Nurses perceived preparedness to screen, intervene, and refer women with suspected postpartum depression. *Health & Social Care in the Community*, 26(1), pp.e75-e84.

Bitew T, Hanlon C and Kebede E ; Honikman S ; Onah M N; Fekadu A ;. (2017). Antenatal depressive symptoms and utilisation of delivery and postnatal care: a prospective study in rural Ethiopia. *BMC Pregnancy & Childbirth*, 17(1), pp.206.

Bittner A, Peukert J and Zimmermann C ; Junge-Hoffmeister J ; Parker L S; Stobel-Richter Y ; Weidner K ;. (2014). Early intervention in pregnant women with elevated anxiety and depressive symptoms: efficacy of a cognitive-behavioral group program. *Journal of Perinatal & Neonatal Nursing*, 28(3), pp.185-95.

Bledsoe Sarah E, Wike Traci and Killian-Farrell Candace ; Lombardi Brianna ; Rizo Cynthia ; Bellows Anne-Marie O; Sommers Amy R; Sheely Amanda L;. (2018). Feasibility of treating depression in pregnant adolescents using brief interpersonal psychotherapy. *Social Work in Mental Health*, 16(3), pp.252-265.

Boath E, Bradley E and Anthony P ;. (2004). Users' views of two alternative approaches to the treatment of postnatal depression. *Journal of Reproductive & Infant Psychology*, 22(1), pp.13-24.

Borghini A, Habersaat S and Forcada-Guex M ; Nessi J ; Pierrehumbert B ; Ansermet F ; Muller-Nix C ;. (2014). Effects of an early intervention on maternal post-traumatic stress symptoms and the quality of mother-infant interaction: the case of preterm birth. *Infant Behavior & Development*, 37(4), pp.624-31.

Bose G N. S. C. (2015). Changes in depression status in low socioeconomic perinatal subjects in rural India after supervised physical exercise: A randomized controlled study. *Indian Journal of Psychiatry*, 57(4), pp.412-413.

Bowen A, Bowen R and Butt P ; Rahman K ; Muhajarine N ;. (2012). Patterns of depression and treatment in pregnant and postpartum women. *Canadian Journal of Psychiatry*, 57(3), pp.161-167.

Bowman Jennifer A and Hellier Susan D; Cline Thomas W;. (2015). Sources Impacting Pharmacological Treatment for Anxiety and/or Depression During Pregnancy. *Journal for Nurse Practitioners*, 11(2), pp.184-191.

Boyd R C, Mogul M and O'Hara M W;. (2015). A Pilot Investigation to Enhance Behavioral Health Referral for Perinatal, Low-Income Women With Mood Disorders. *Progress in Community Health Partnerships*, 9(4), pp.583-9.

Boyd R C, Price J and Mogul M ; Yates T ; Guevara J P;. (2019). Pilot RCT of a social media parenting intervention for postpartum mothers with depression symptoms. *Journal of Reproductive & Infant Psychology*, 37(3), pp.290-301.

Brandon A R, Ceccotti N and Hynan L S; Shivakumar G ; Johnson N ; Jarrett R B;. (2012). Proof of concept: Partner-Assisted Interpersonal Psychotherapy for perinatal depression. *Archives of Women's Mental Health*, 15(6), pp.469-80.

Brecht Carrie Jean. (2014). Prevention of postpartum traumatic stress in mothers with preterm infants: Manual development and evaluation. *Dissertation Abstracts International: Section B: The Sciences and Engineering*, 74(11-B(E)), pp.No Pagination Specified.

Brock R L and O'Hara M W; Segre L S;. (2017). Depression Treatment by Non-Mental-Health Providers: Incremental Evidence for the Effectiveness of Listening Visits. *American Journal of Community Psychology*, 59(1-2), pp.172-183.

Broom M A and Ladley A S; Rhyne E A; Halloran D R;. (2015). Feasibility and Perception of Using Text Messages as an Adjunct Therapy for Low-Income, Minority Mothers With Postpartum Depression. *JMIR Mental Health*, 2(1), pp.e4.

Brugha T S, Wheatley S and Taub N A; Culverwell A ; Friedman T ; Kirwan P ; Jones D R; Shapiro D A;. (2000). Pragmatic randomized trial of antenatal intervention to prevent postnatal depression by reducing psychosocial risk factors. *Psychological Medicine*, 30(6), pp.1273-81.

Brugha T S, Smith J and Austin J ; Bankart J ; Patterson M ; Lovett C ; Morgan Z ; Morrell C J; Slade P ;. (2016). Can community midwives prevent antenatal depression? An external

pilot study to test the feasibility of a cluster randomized controlled universal prevention trial. *Psychological Medicine*, 46(2), pp.345-56.

Buil Aude, Caeymaex Laurence and Mero Sophie ; Sankey Carol ; Apter Gisèle ; Devouche Emmanuel ;. (2019). Kangaroo supported diagonal flexion positioning: Positive impact on maternal stress and postpartum depression risk and on skin-to-skin practice with very preterm infants. *Journal of Neonatal Nursing*, 25(2), pp.86-92.

Buist A, Condon J and Brooks J ; Speelman C ; Milgrom J ; Hayes B ; Ellwood D ; Barnett B ; Kowalenko N ; Matthey S ; Austin M P; Bilszta J ;. (2006). Acceptability of routine screening for perinatal depression. *Journal of Affective Disorders*, 93(1-3), pp.233-7.

Buist A, Speelman C and Hayes B ; Reay R ; Milgrom J ; Meyer D ; Condon J ;. (2007). Impact of education on women with perinatal depression. *Journal of Psychosomatic Obstetrics & Gynecology*, 28(1), pp.49-54.

Buist A, Ellwood D and Brooks J ; Milgrom J ; Hayes B A; Sved-Williams A ; Barnett B ; Karatas J ; Bilszta J ;. (2007). National program for depression associated with childbirth: the Australian experience. *Best Practice & Research in Clinical Obstetrics & Gynaecology*, 21(2), pp.193-206.

Burns A, O'Mahen H and Baxter H ; Bennert K ; Wiles N ; Ramchandani P ; Turner K ; Sharp D ; Thorn J ; Noble S ; Evans J ;. (2013). A pilot randomised controlled trial of cognitive behavioural therapy for antenatal depression. *BMC Psychiatry*, 13, pp.33.

Butler Hannah, Hare Dougal and Walker Samantha ; Wieck Angelika ; Wittkowski Anja ;. (2014). The acceptability and feasibility of the Baby Triple P Positive Parenting Programme on a mother and baby unit: Q-methodology with mothers with severe mental illness. *Archives of Women's Mental Health*, 17(5), pp.455-463.

Buttner Melissa Mercedes. (2014). Efficacy of yoga for depressed postpartum women: A randomized controlled trial. *Dissertation Abstracts International: Section B: The Sciences and Engineering*, 75(1-B(E)), pp.No Pagination Specified.

Buultjens M, Robinson P and Liamputtong P ;. (2008). A holistic programme for mothers with postnatal depression: pilot study. *Journal of Advanced Nursing*, 63(2), pp.181-8.

Buzi R S and Smith P B; Kozinetz C A; Peskin M F; Wiemann C M;. (2015). A Socioecological Framework to Assessing Depression Among Pregnant Teens. *Maternal & Child Health Journal*, 19(10), pp.2187-94.

Byatt N, Pbert L and Hosein S ; Swartz H A; Weinreb L ; Allison J ; Ziedonis D ;. (2016). PRogram in Support of Moms (PRISM): Development and beta testing. *Psychiatric Services*, 67(8), pp.824-826.

Byatt N, Biebel K and Moore Simas ; T A ; Sarvet B ; Ravech M ; Allison J ; Straus J ;. (2016). Improving perinatal depression care: the Massachusetts Child Psychiatry Access Project for Moms. *General Hospital Psychiatry*, 40, pp.12-7.

Byatt N, Straus J and Stopa A ; Biebel K ; Mittal L ; Moore Simas ; T A ;. (2018). Massachusetts Child Psychiatry Access Program for Moms: Utilization and Quality Assessment. *Obstetrics & Gynecology*, 132(2), pp.345-353.



Byatt N, Moore Simas and T A ; Biebel K ; Sankaran P ; Pbert L ; Weinreb L ; Ziedonis D ; Allison J ;. (2018). PRogram In Support of Moms (PRISM): a pilot group randomized controlled trial of two approaches to improving depression among perinatal women. *Journal of Psychosomatic Obstetrics & Gynecology*, 39(4), pp.297-306.

Callahan J L and Hynan M T;. (2002). Identifying mothers at risk for postnatal emotional distress: further evidence for the validity of the Perinatal Posttraumatic Stress Disorder Questionnaire. *Journal of Perinatology*, 22(6), pp.448-454.

Calvert Cathryn A. (2009). Structural family therapy as a treatment modality to decrease depressive symptoms for women suffering from postpartum depression and improve family functioning. *Dissertation Abstracts International: Section B: The Sciences and Engineering*, 69(7-B), pp.4079.

Cardone I A and Kim J J; Gordon T E; Gordon S M; Silver R K;. (2006). Psychosocial assessment by phone for high-scoring patients taking the Edinburgh Postnatal Depression Scale: communication pathways and strategies. *Archives of Women's Mental Health*, 9(2), pp.87-94.

Carlin E, Atkinson D and Marley J V;. (2019). 'Having a Quiet Word': Yarning with Aboriginal Women in the Pilbara Region of Western Australia about Mental Health and Mental Health Screening during the Perinatal Period. *International Journal of Environmental Research & Public Health [Electronic Resource]*, 16(21), pp.01.

Carolan Mary, Barry Maebh and Gamble Mary ; Turner Kathleen ; Mascareñas Óscar ;. (2012). Experiences of pregnant women attending a lullaby programme in Limerick, Ireland: A qualitative study. *Midwifery*, 28(3), pp.321-328.

Carolan M, Barry M and Gamble M ; Turner K ; Mascarenas O ;. (2012). The Limerick Lullaby project: an intervention to relieve prenatal stress. *Midwifery*, 28(2), pp.173-80.

Carter R, Cust F and Boath E ;. (2019). Effectiveness of a peer support intervention for antenatal depression: a feasibility study. *Journal of Reproductive & Infant Psychology*, , pp.1-12.

Chae S Y, Chae M H; Tyndall A and Ramirez M R; Winter R O;. (2012). Can we effectively use the two-item PHQ-2 to screen for postpartum depression?. *Family Medicine*, 44(10), pp.698-703.

Chambers Andrea S. (2009). Relaxation during pregnancy to reduce stress and anxiety and their associated complications. *Dissertation Abstracts International: Section B: The Sciences and Engineering*, 69(7-B), pp.4412.

Chang M Y and Chen C H; Huang K F;. (2008). Effects of music therapy on psychological health of women during pregnancy. *Journal of Clinical Nursing*, 17(19), pp.2580-7.

Chang H C and Yu C H; Chen S Y; Chen C H;. (2015). The effects of music listening on psychosocial stress and maternal-fetal attachment during pregnancy. *Complementary Therapies in Medicine*, 23(4), pp.509-15.

Chaudhry Simriti K and Gordon-Elliott Janna S; Brody Benjamin D;. (2016). The Cornell Peripartum Psychosis Management Tool: A case series and template. *Psychosomatics: Journal of Consultation and Liaison Psychiatry*, 57(3), pp.319-324.

Chaudron L H and Szilagyi P G; Kitzman H J; Wadkins H I; Conwell Y. (2004). Detection of postpartum depressive symptoms by screening at well-child visits. *Pediatrics*, 113(3 Pt 1), pp.551-8.

Chen C H and Tseng Y F; Chou F H; Wang S Y;. (2000). Effects of support group intervention in postnatally distressed women. A controlled study in Taiwan. *Journal of Psychosomatic Research*, 49(6), pp.395-9.

Chen H, Wang J and Ch'ng Y C; Mingoo R ; Lee T ; Ong J ;. (2011). Identifying mothers with postpartum depression early: integrating perinatal mental health care into the obstetric setting. *ISRN Obstetrics & Gynecology*, 2011, pp.309189.

Chen Helen. (2011). Addressing maternal mental health needs in Singapore. *Psychiatric Services*, 62(1), pp.102.

Cheng Erika R, Rifas-Shiman Sheryl L; Perkins Meghan E; Rich-Edwards Janet Wilson; Gillman Matthew W; Wright Rosalind and Taveras Elsie M;. (2016). The Influence of Antenatal Partner Support on Pregnancy Outcomes. *Journal of Women's Health (15409996)*, 25(7), pp.672-679.

Cherry A S, Blucker R T; Thornberry T S; Hetherington C and McCaffree M A; Gillaspay S R;. (2016). Postpartum depression screening in the Neonatal Intensive Care Unit: program development, implementation, and lessons learned. *Journal of multidisciplinary healthcare*, 9, pp.59-67.

Chibanda D, Shetty A K and Tshimanga M ; Woelk G ; Stranix-Chibanda L ; Rusakaniko S ;. (2014). Group problem-solving therapy for postnatal depression among HIV-positive and HIV-negative mothers in Zimbabwe. *Journal of the International Association of Providers of AIDS Care*, 13(4), pp.335-41.

Christie J and Bunting B . (2011). The effect of health visitors' postpartum home visit frequency on first-time mothers: cluster randomised trial. *International Journal of Nursing Studies*, 48(6), pp.689-702.

Chuffo Siewert, R and Cline M ; Segre L S;. (2015). Implementation of an innovative nurse-delivered depression intervention for mothers of NICU infants. *Advances in Neonatal Care*, 15(2), pp.104-11.

Chung K F, Yeung W F; Zhang Z J; Yung K P; Man S C; Lee C P; Lam S K; Leung T W; Leung K Y; Ziea E T; Taam Wong and V ;. (2012). Randomized non-invasive sham-controlled pilot trial of electroacupuncture for postpartum depression. *Journal of Affective Disorders*, 142(1-3), pp.115-21.

Clarici A, Pellizzoni S and Guaschino S ; Alberico S ; Bembich S ; Giuliani R ; Short A ; Guarino G ; Panksepp J ;. (2015). Intranasal administration of oxytocin in postnatal depression: implications for psychodynamic psychotherapy from a randomized double-blind pilot study. *Frontiers in Psychology*, 6, pp.426.

Clark G. (2000). Discussing emotional health in pregnancy: the Edinburgh Postnatal Depression Scale. *British Journal of Community Nursing*, 5(2), pp.91-8.

Clarke K, Azad K and Kuddus A ; Shaha S ; Nahar T ; Aumon B H; Hossen M M; Beard J ; Costello A ; Houweling T A; Prost A ; Fottrell E ;. (2014). Impact of a participatory

intervention with women's groups on psychological distress among mothers in rural Bangladesh: secondary analysis of a cluster-randomised controlled trial. *PLoS ONE [Electronic Resource]*, 9(10), pp.e110697.

Clevesy M A, Gatlin T K; Cheese C and Strebel K ;. (2019). A Project to Improve Postpartum Depression Screening Practices Among Providers in a Community Women's Health Care Clinic. *Nursing for Women's Health*, 23(1), pp.21-30.

Cluxton-Keller Fallon, Buteau Jennifer and Williams Melony ; Stolte Patricia ; Monroe-Cassel Maggie ; Bruce Martha ;. (2019). Engaging rural young mothers in a technology-based intervention for depression. *Child & Youth Services*, , pp.No Pagination Specified.

Coates D, Davis E and Campbell L ;. (2017). The experiences of women who have accessed a perinatal and infant mental health service: A qualitative investigation. *Advances in Mental Health*, 15(1), pp.88-100.

Cobiella C W and Mabe P A; Forehand R L;. (1990). A comparison of two stress-reduction treatments for mothers of neonates hospitalized in a neonatal intensive care unit. *Children's Health Care*, 19(2), pp.93-100.

Coe C and Barlow J . (2013). Supporting women with perinatal mental health problems: the role of the voluntary sector. *Community Practitioner*, 86(2), pp.23-7.

Coffman M J, Scott V and Schuch C ; Mele C ; Mayfield C ; Balasubramanian V ; Stevens A ; Dulin M ;. (2019). Postpartum Depression Screening and Referrals in Special Supplemental Nutrition Program for Women, Infant, and Children Clinics. *JOGNN Journal of Obstetric, Gynecologic and & Neonatal Nursing*, 29, pp.29.

Cohen J I. (2002). *Effects of an intervention program to decrease distress in pregnant Latina women at risk for preterm birth*. . University of California, Los Angeles.

Cole J C. M, Olkkola M and Zarrin H E; Berger K ; Moldenhauer J S;. (2018). Universal Postpartum Mental Health Screening for Parents of Newborns With Prenatally Diagnosed Birth Defects. *JOGNN - Journal of Obstetric, Gynecologic and & Neonatal Nursing*, 47(1), pp.84-93.

Coll C V. N, Domingues M R; Stein A and da Silva ; B G C; Bassani D G; Hartwig F P; da Silva ; I C M; da Silveira ; M F ; da Silva ; S G ; Bertoldi A D;. (2019). Efficacy of Regular Exercise During Pregnancy on the Prevention of Postpartum Depression: The PAMELA Randomized Clinical Trial. *JAMA network open*, 2(1), pp.e186861.

Cooper P J, De Pascalis and L ; Woolgar M ; Romaniuk H ; Murray L ;. (2015). Attempting to prevent postnatal depression by targeting the mother-infant relationship: a randomised controlled trial. *Primary Health Care Research & Development*, 16(4), pp.383-97.

Corey K, Fallek R and Benattar M ;. (2019). Bedside Music Therapy for Women during Antepartum and Postpartum Hospitalization. *MCN and American Journal of Maternal Child Nursing*, 44(5), pp.277-283.

Corna Francesca, Tofail Fahmida and Chowdhury Mita Rani Roy; Bizouerne Cécile ;. (2019). Supporting maternal mental health of Rohingya refugee women during the perinatal period to promote child health and wellbeing: a field study in Cox's Bazar. *Intervention (15718883)*, 17(2), pp.160-168.

Corno G, Etchemendy E and Espinoza M ; Herrero R ; Molinari G ; Carrillo A ; Drossaert C ; Banos R M;. (2018). Effect of a web-based positive psychology intervention on prenatal well-being: A case series study. *Women and Birth*, 31(1), pp.e1-e8.

Craig E, Judd F and Hodgins G ;. (2005). Therapeutic group programme for women with postnatal depression in rural Victoria: a pilot study. *Australasian Psychiatry*, 13(3), pp.291-5.

Crockett K, Zlotnick C and Davis M ; Payne N ; Washington R ;. (2008). A depression preventive intervention for rural low-income African-American pregnant women at risk for postpartum depression. *Archives of Women's Mental Health*, 11(5-6), pp.319-25.

Curry M A, Burton D and Fields J ;. (1998). The Prenatal Psychosocial Profile: a research and clinical tool. *Research in Nursing & Health*, 21(3), pp.211-219.

Cussino M, Zaccagnino M and Fini G D. I; Veglia F ;. (2013). Postnatal depression and prevention: Effects of a psychoeducational intervention. [Italian, English]. *Minerva Psichiatrica*, 54(1), pp.41-58.

Cwikel Julie, Segal-Engelchin Dorit and Niego Limor ;. (2018). Addressing the needs of new mothers in a multi-cultural setting: an evaluation of home visiting support for new mothers - Mom to Mom (Negev). *Psychology and Health & Medicine*, 23(5), pp.517-524.

Cynthia Logsdon, M and Myers J ; Rushton J ; Gregg J L; Josephson A M; Davis D W; Brothers K ; Baisch K ; Carabello A ; Vogt K ; Jones K ; Angermeier J ;. (2018). Efficacy of an Internet-based depression intervention to improve rates of treatment in adolescent mothers. *Archives of Women's Mental Health*, 21(3), pp.273-285.

Da Costa , D and Lowensteyn I ; Abrahamowicz M ; Ionescu-Iltu R ; Dritsa M ; Rippen N ; Cervantes P ; Khalife S ;. (2009). A randomized clinical trial of exercise to alleviate postpartum depressed mood. *Journal of Psychosomatic Obstetrics & Gynecology*, 30(3), pp.191-200.

Dabas Suman, Joshi Poonam and Agarwal Ramesh ; Yadav Raj Kumar; Kachhawa Garima ;. (2019). Impact of audio assisted relaxation technique on stress, anxiety and milk output among postpartum mothers of hospitalized neonates: A randomized controlled trial. *Journal of Neonatal Nursing*, 25(4), pp.200-204.

Dagher R K and McGovern P M; Alexander B H; Dowd B E; Ukestad L K; McCaffrey D J;. (2009). The psychosocial work environment and maternal postpartum depression. *International Journal of Behavioral Medicine*, 16(4), pp.339-46.

Dagklis T, Papazisis G and Tsakiridis I ; Chouliara F ; Mamopoulos A ; Rousso D ;. (2016). Prevalence of antenatal depression and associated factors among pregnant women hospitalized in a high-risk pregnancy unit in Greece. *Social Psychiatry & Psychiatric Epidemiology*, 51(7), pp.1025-31.

Daire Andrew P, Liu Xun and Tucker Kelsee ; Williams Brooke ; Broyles Amanda ; Wheeler Naomi ;. (2018). Positively impacting maternal stress and parental adjustment through community-based relationship education (re). *Marriage & Family Review*, , pp.No Pagination Specified.

Daley A, Winter H and Grimmett C ; McGuinness M ; McManus R ; MacArthur C ;. (2008). Feasibility of an exercise intervention for women with postnatal depression: a pilot randomised controlled trial. *British Journal of General Practice*, 58(548), pp.178-83.

Daley A J, Blamey R V; Jolly K and Roalfe A K; Turner K M; Coleman S ; McGuinness M ; Jones I ; Sharp D J; MacArthur C ;. (2015). A pragmatic randomized controlled trial to evaluate the effectiveness of a facilitated exercise intervention as a treatment for postnatal depression: the PAM-PeRS trial. *Psychological Medicine*, 45(11), pp.2413-25.

Danaher B G, Milgrom J and Seeley J R; Stuart S ; Schembri C ; Tyler M S; Ericksen J ; Lester W ; Gemmill A W; Lewinsohn P ;. (2012). Web-Based Intervention for Postpartum Depression: Formative Research and Design of the MomMoodBooster Program. *JMIR Research Protocols*, 1(2), pp.e18.

Danaher B G, Milgrom J and Seeley J R; Stuart S ; Schembri C ; Tyler M S; Ericksen J ; Lester W ; Gemmill A W; Kosty D B; Lewinsohn P ;. (2013). MomMoodBooster web-based intervention for postpartum depression: feasibility trial results. *Journal of Medical Internet Research*, 15(11), pp.e242.

Darius Tandon, S and Leis J A; Ward E A; Snyder H ; Mendelson T ; Perry D F; Carter M ; Hamil J ; Le H N ;. (2018). Adaptation of an evidence-based postpartum depression intervention: feasibility and acceptability of mothers and babies 1-on-1. *BMC Pregnancy & Childbirth*, 18(1), pp.93.

Darwin Z, McGowan L and Edozien L C;. (2013). Assessment acting as intervention: Findings from a study of perinatal psychosocial assessment. *Journal of Reproductive and Infant Psychology*, 31(5), pp.500-511.

Davies Simone, Horne Rebecca and Moore Tal ;. (2018). Psychological Interventions in Perinatal Community Mental Health Teams in the United Kingdom. *Journal of Prenatal & Perinatal Psychology & Health*, 33(1), pp.27-37.

Davis Kyle J. (2014). The feasibility of yoga in the treatment of antenatal depression and anxiety: A pilot study. *Dissertation Abstracts International: Section B: The Sciences and Engineering*, 74(12-B(E)), pp.No Pagination Specified.

Davis K, Goodman S H and Leiferman J ; Taylor M ; Dimidjian S ;. (2015). A randomized controlled trial of yoga for pregnant women with symptoms of depression and anxiety. *Complementary Therapies in Clinical Practice*, 21(3), pp.166-72.

de Alencar , A E and Arraes L C; de Albuquerque ; E C ; Alves J G;. (2009). Effect of kangaroo mother care on postpartum depression. *Journal of Tropical Pediatrics*, 55(1), pp.36-8.

de la Fe Rodriguez-Munoz, M and Le H N ; de la Cruz ; I V ; Crespo M E. O; Mendez N I;. (2017). Feasibility of screening and prevalence of prenatal depression in an obstetric setting in Spain. *European Journal of Obstetrics, Gynecology and & Reproductive Biology*, 215, pp.101-105.

De Schepper , S and Vercauteren T ; Tersago J ; Jacquemyn Y ; Raes F ; Franck E ;. (2016). Post-Traumatic Stress Disorder after childbirth and the influence of maternity team care during labour and birth: A cohort study. *Midwifery*, 32, pp.87-92.

Dennis C L. (2003). The effect of peer support on postpartum depression: a pilot randomized controlled trial. *Canadian Journal of Psychiatry - Revue Canadienne de Psychiatrie*, 48(2), pp.115-24.

Dennis C L. (2004). Can we identify mothers at risk for postpartum depression in the immediate postpartum period using the Edinburgh Postnatal Depression Scale?. *Journal of Affective Disorders*, 78(2), pp.163-9.

Dennis C L. (2010). Postpartum depression peer support: maternal perceptions from a randomized controlled trial. *International Journal of Nursing Studies*, 47(5), pp.560-8.

Dennis C L. (2013). Peer support for postpartum depression: volunteers' perceptions, recruitment strategies and training from a randomized controlled trial. *Health Promotion International*, 28(2), pp.187-96.

Dennis C L. (2014). The process of developing and implementing a telephone-based peer support program for postpartum depression: evidence from two randomized controlled trials. *Trials [Electronic Resource]*, 15, pp.131.

Dennis-Tiway T A, Denefrio S and Gelber S ;. (2017). Salutary effects of an attention bias modification mobile application on biobehavioral measures of stress and anxiety during pregnancy. *Biological Psychology*, 127, pp.148-156.

DeOliveira Cherish A. (2018). A screening tool for perinatal mood and anxiety disorders: Culturally sensitive prenatal care for Latina mothers. *Dissertation Abstracts International: Section B: The Sciences and Engineering*, 79(2-B(E)), pp.No Pagination Specified.

Deshpande C S, Rakhshani A and Nagarathna R ; Ganpat T S; Kurpad A ; Maskar R ; Nagendra H R; Sudheer D C; Abbas R ; Raghuram N ; Anura K ; Rita M ; Ramarao N ;. (2013). Yoga for high-risk pregnancy: a randomized controlled trial. *Annals of Medical & Health Sciences Research*, 3(3), pp.341-4.

Dikmen-Yildiz P, Ayers S and Phillips L ;. (2017). Screening for birth-related PTSD: psychometric properties of the Turkish version of the Posttraumatic Diagnostic Scale in postpartum women in Turkey. *European Journal of Psychotraumatology*, 8(1), pp.1306414.

Dimidjian S, Goodman S H and Felder J N; Gallop R ; Brown A P; Beck A ;. (2015). An open trial of mindfulness-based cognitive therapy for the prevention of perinatal depressive relapse/recurrence. *Archives of Women's Mental Health*, 18(1), pp.85-94.

Dimidjian S, Goodman S H and Felder J N; Gallop R ; Brown A P; Beck A ;. (2016). Staying well during pregnancy and the postpartum: A pilot randomized trial of mindfulness-based cognitive therapy for the prevention of depressive relapse/recurrence. *Journal of Consulting & Clinical Psychology*, 84(2), pp.134-45.

Ding T, Wang D X and Qu Y ; Chen Q ; Zhu S N;. (2014). Epidural labor analgesia is associated with a decreased risk of postpartum depression: a prospective cohort study. *Anesthesia & Analgesia*, 119(2), pp.383-92.

Docherty A, Najjar R and Combs S ; Woolley R ; Stoyles S ;. (2019). Postpartum depression screening in the first year: A cross-sectional provider analysis in Oregon. *Journal of the American Association of Nurse Practitioners*, 31, pp.31.

Dodge K A, Goodman W B; Murphy R A; O'Donnell K and Sato J ; Guptill S ;. (2014). Implementation and randomized controlled trial evaluation of universal postnatal nurse home visiting. *American Journal of Public Health*, 104 Suppl 1, pp.S136-43.

Doherty K, Barry M and Marcano-Belisario J ; Arnaud B ; Morrison C ; Car J ; Doherty G ;. (2018). A Mobile App for the Self-Report of Psychological Well-Being During Pregnancy (BrightSelf): Qualitative Design Study. *JMIR Mental Health*, 5(4), pp.e10007.

Dois C A and Lucchini R C; Villarroel D L; Uribe T C;. (2013). The effect of Mother/Infant skin-to-skin contact on postpartum depressive symptoms in women with low obstetric risk. *Revista Chilena de Pediatría*, 84(3), pp.285-292.

Dolan R, Hann M and Edge D ; Shaw J ;. (2019). Pregnancy in prison, mental health and admission to prison mother and baby units. *Journal of Forensic Psychiatry and Psychology*, 30(3), pp.448-466.

Dombrowska-Pali A, Gebuza G and Kazmierczak M ; Gierszewska M ; Mieczkowska E ;. (2018). Music therapy - An alternative form supporting the quality of mother and baby care. *Pediatrica Polska*, 93(4), pp.331-335.

Dornelas E A, Ferrand J and Stepnowski R ; Barbagallo J ; McCullough L ;. (2010). A Pilot Study of Affect-Focused Psychotherapy for Antepartum Depression. *Journal of Psychotherapy Integration*, 20(4), pp.364-382.

Downie J, Wynaden D and McGowan S ; Juliff D ; Axten C ; Fitzpatrick L ; Ogilvie S ; Painter S ;. (2003). Using the Edinburgh Postnatal Depression Scale to achieve best practice standards. *Nursing & Health Sciences*, 5(4), pp.283-7.

Drake E, Howard E and Kinsey E ;. (2014). Online screening and referral for postpartum depression: an exploratory study. *Community Mental Health Journal*, 50(3), pp.305-11.

Drozd F, Haga S M and Brendryen H ; Slinning K ;. (2015). An Internet-Based Intervention (Mamma Mia) for Postpartum Depression: Mapping the Development from Theory to Practice. *JMIR Research Protocols*, 4(4), pp.e120.

Dubus Nicole. (2014). Permission to be authentic: An intervention for postpartum women. *Affilia: Journal of Women & Social Work*, 29(1), pp.43-55.

Duffecy J, Grekin R and Hinkel H ; Gallivan N ; Nelson G ; O'Hara M W;. (2019). A Group-Based Online Intervention to Prevent Postpartum Depression (Sunnyside): Feasibility Randomized Controlled Trial. *JMIR Mental Health*, 6(5), pp.e10778.

Dugravier R, Tubach F and Saias T ; Guedeney N ; Pasquet B ; Purper-Ouakil D ; Tereno S ; Welniarz B ; Matos J ; Group Capedp Study; Guedeney A ; Greacen T ;. (2013). Impact of a manualized multifocal perinatal home-visiting program using psychologists on postnatal depression: the CAPEDP randomized controlled trial. *PLoS ONE [Electronic Resource]*, 8(8), pp.e72216.

Edge Dawn. (2011). Managed care networks improve perinatal mental healthcare for Black and minority ethnic (BME) women?. *Journal of Public Mental Health*, 10(3), pp.151-163.

- Edward K L, Giandinoto J A; Stephenson J and Mills C ; McFarland J ; Castle D J;. (2019). Self- screening using the Edinburgh post natal depression scale for mothers and fathers to initiate early help seeking behaviours. *Archives of Psychiatric Nursing*, 33(4), pp.421-427.
- Edwards B, Galletly C and Semmler-Booth T ; Dekker G ;. (2008). Antenatal psychosocial risk factors and depression among women living in socioeconomically disadvantaged suburbs in Adelaide, South Australia. *Australian & New Zealand Journal of Psychiatry*, 42(1), pp.45-50.
- Lonnberg G, Jonas W and Unternaehrer E ; Branstrom R ; Nissen E ; Niemi M ;. ( ). Effects of a mindfulness based childbirth and parenting program on pregnant women's perceived stress and risk of perinatal depression-Results from a randomized controlled trial. *Journal of Affective Disorders*, 262, pp.133-142.
- El-Khorazaty M N, Johnson A A; Kiely M and El-Mohandes A A; Subramanian S ; Laryea H A; Murray K B; Thornberry J S; Joseph J G;. (2007). Recruitment and retention of low-income minority women in a behavioral intervention to reduce smoking, depression, and intimate partner violence during pregnancy. *BMC Public Health*, 7, pp.233.
- Elliott S A, Gerrard J and Ashton C ; Cox J L;. (2001). Training health visitors to reduce levels of depression after childbirth: an evaluation. *Journal of Mental Health*, 10(6), pp.613-625.
- Emerson M R and Mathews T L; Struwe L. (2018). Postpartum Depression Screening for New Mothers at Well Child Visits. *MCN and American Journal of Maternal Child Nursing*, 43(3), pp.139-145.
- Engqvist I and Nilsson K . (2013). Involving the family in the care and treatment of women with postpartum psychosis: Swedish psychiatrists' experiences. *Psychiatry Journal Print*, 2013, pp.897084.
- Ettenberger M, Beltran Ardila and Y M ;. (2018). Music therapy song writing with mothers of preterm babies in the Neonatal Intensive Care Unit (NICU) - A mixed-methods pilot study. *Arts in Psychotherapy*, 58, pp.42-52.
- Evans Amanda P, Goodman Sherryl H; Dimidjian Sona and Gallop Robert ;. (2019). The role of engagement in mindfulness-based cognitive therapy for the prevention of depressive relapse/recurrence in perinatal women. *Mindfulness*, , pp.No Pagination Specified.
- Falcone V M, Mader C V; Nascimento C F; Santos J M; de Nobrega and F J ;. (2005). [Multiprofessional care and mental health in pregnant women]. *Revista de Saude Publica*, 39(4), pp.612-8.
- Fancourt D and Perkins R . (2019). Creative interventions for symptoms of postnatal depression: A process evaluation of implementation. *Arts & Health*, 11(1), pp.38-53.
- Feeley N, Zekowitz P and Charbonneau L ; Cormier C ; Lacroix A ; Marie C S; Papageorgiou A ;. (2008). Assessing the feasibility and acceptability of an intervention to reduce anxiety and enhance sensitivity among mothers of very low birth-weight infants. *Advances in Neonatal Care*, 8(5), pp.276-84.



Feinberg E, Stein R and Diaz-Linhart Y ; Egbert L ; Beardslee W ; Hegel M T; Silverstein M ;. (2012). Adaptation of problem-solving treatment for prevention of depression among low-income, culturally diverse mothers. *Family & Community Health*, 35(1), pp.57-67.

Felder Jennifer Nicole. (2016). Targeting barriers to care for pregnant women at risk for depression: Examining the role of stigma and the feasibility of a web-based depression prevention program. *Dissertation Abstracts International: Section B: The Sciences and Engineering*, 77(1-B(E)), pp.No Pagination Specified.

Felder J N, Segal Z and Beck A ; Sherwood N E; Goodman S H; Boggs J ; Lemon E ; Dimidjian S ;. (2017). An Open Trial of Web-Based Mindfulness-Based Cognitive Therapy for Perinatal Women at Risk for Depressive Relapse. *Cognitive and Behavioral Practice*, 24(1), pp.26-37.

Felder J N, Epel E and Lewis J B; Cunningham S D; Tobin J N; Rising S S; Thomas M ; Ickovics J R;. (2017). Depressive symptoms and gestational length among pregnant adolescents: Cluster randomized control trial of CenteringPregnancy plus group prenatal care. *Journal of Consulting & Clinical Psychology*, 85(6), pp.574-584.

Felder J N, Laraia B and Coleman-Phox K ; Bush N ; Suresh M ; Thomas M ; Adler N ; Epel E ; Prather A A;. (2018). Poor Sleep Quality, Psychological Distress, and the Buffering Effect of Mindfulness Training During Pregnancy. *Behavioral Sleep Medicine*, 16(6), pp.611-624.

Figueira P, Correa H and Malloy-Diniz L ; Romano-Silva M A;. (2009). Edinburgh Postnatal Depression Scale for screening in the public health system. *Revista de Saude Publica*, 43 Suppl 1, pp.79-84.

Figueroa-Leigh F, Rojas P and Castanon C ;. (2015). Screening for postpartum depression in a private health care network in Chile. *Family Practice*, 32(4), pp.431-5.

Filoche Sara, Lawton Bev and Stanley James ;. (2016). Access to Secondary Mental Health Services in a Cohort of New Zealand Mothers. *Community Mental Health Journal*, 52(8), pp.964-971.

Fisher J, Astbury J and Smith A ;. (1997). Adverse psychological impact of operative obstetric interventions: a prospective longitudinal study. *Australian & New Zealand Journal of Psychiatry*, 31(5), pp.728-38.

Fisher J R and Wynter K H; Rowe H J;. (2010). Innovative psycho-educational program to prevent common postpartum mental disorders in primiparous women: a before and after controlled study. *BMC Public Health*, 10, pp.432.

Fisher J, Chatham E and Haseler S ; McGaw B ; Thompson J ;. (2012). Uneven implementation of the National Perinatal Depression Initiative: findings from a survey of Australian women's hospitals. *Australian & New Zealand Journal of Obstetrics & Gynaecology*, 52(6), pp.559-64.

Fisher J, Rowe H and Wynter K ; Tran T ; Lorgelly P ; Amir L H; Proimos J ; Ranasinha S ; Hiscock H ; Bayer J ; Cann W ;. (2016). Gender-informed, psychoeducational programme for couples to prevent postnatal common mental disorders among primiparous women: cluster randomised controlled trial. *BMJ Open*, 6(3), pp.e009396.

Flanagan T and Avalos L A. (2016). Perinatal Obstetric Office Depression Screening and Treatment: Implementation in a Health Care System. *Obstetrics & Gynecology*, 127(5), pp.911-5.

Flynn H A, O'Mahen H A; Massey L and Marcus S ;. (2006). The impact of a brief obstetrics clinic-based intervention on treatment use for perinatal depression. *Journal of Women's Health*, 15(10), pp.1195-204.

Flynn H A, Sexton M and Ratliff S ; Porter K ; Zivin K ;. (2011). Comparative performance of the Edinburgh Postnatal Depression Scale and the Patient Health Questionnaire-9 in pregnant and postpartum women seeking psychiatric services. *Psychiatry Research*, 187(1-2), pp.130-4.

Flynn Theresa, Jones Brittney A and Ausderau Karla K;. (2016). The Effects of Guided Imagery on Stress in Pregnant Adolescents. *American Journal of Occupational Therapy*, 70, pp.1-1.

Fonseca A, Gorayeb R and Canavarro M C;. (2016). Women's use of online resources and acceptance of e-mental health tools during the perinatal period. *International Journal of Medical Informatics*, 94, pp.228-36.

Fonseca A, Pereira M and Araujo-Pedrosa A ; Gorayeb R ; Ramos M M; Canavarro M C;. (2018). Be a Mom: Formative Evaluation of a Web-Based Psychological Intervention to Prevent Postpartum Depression. *Cognitive and Behavioral Practice*, 25(4), pp.473-495.

Forsell E, Bendix M and Hollandare F ; Szymanska von Schultz; B ; Nasiell J ; Blomdahl-Wetterholm M ; Eriksson C ; Kvarned S ; Lindau van der Linden; J ; Soderberg E ; Jokinen J ; Wide K ; Kaldo V ;. (2017). Internet delivered cognitive behavior therapy for antenatal depression: A randomised controlled trial. *Journal of Affective Disorders*, 221, pp.56-64.

Forsyth J, Boath E and Henshaw C ; Brown H ;. (2017). Exercise as an adjunct treatment for postpartum depression for women living in an inner city-A pilot study. *Health care for women international*, 38(6), pp.635-639.

Fossey L, Papiernik E and Bydlowski M ;. (1997). Postpartum blues: a clinical syndrome and predictor of postnatal depression?. *Journal of Psychosomatic Obstetrics & Gynecology*, 18(1), pp.17-21.

Freeman M P, Wright R and Watchman M ; Wahl R A; Sisk D J; Fraleigh L ; Weibrecht J M;. (2005). Postpartum depression assessments at well-baby visits: screening feasibility, prevalence, and risk factors. *Journal of Women's Health*, 14(10), pp.929-35.

Freeman M P, Davis M and Sinha P ; Wisner K L; Hibbeln J R; Gelenberg A J;. (2008). Omega-3 fatty acids and supportive psychotherapy for perinatal depression: a randomized placebo-controlled study. *Journal of Affective Disorders*, 110(1-2), pp.142-8.

Fuggle P and Haydon K . (2000). Estimating resources for setting up an HV service for postnatal depression. *British Journal of Community Nursing*, 5(7), pp.348-51.

Fuhr D C, Weobong B and Lazarus A ; Vanobberghen F ; Weiss H A; Singla D R; Tabana H ; Afonso E ; De Sa ; A ; D'Souza E ; Joshi A ; Korgaonkar P ; Krishna R ; Price L N; Rahman A ; Patel V ;. (2019). Delivering the Thinking Healthy Programme for perinatal depression

through peers: an individually randomised controlled trial in India. *The Lancet Psychiatry*, 6(2), pp.115-127.

Gamble J, Creedy D and Moyle W ;. (2004). Counselling processes to address psychological distress following childbirth: perceptions of women. *Australian Midwifery*, 17(3), pp.12-15.

Gamble J, Creedy D and Moyle W ;. (2004). Counselling processes to address psychological distress following childbirth: perceptions of midwives. *Australian Midwifery*, 17(3), pp.16-19.

Gao L L and Luo S Y; Chan S W;. (2012). Interpersonal psychotherapy-oriented program for Chinese pregnant women: delivery, content, and personal impact. *Nursing & Health Sciences*, 14(3), pp.318-24.

Gao L L and Chan S W; Sun K. (2012). Effects of an interpersonal-psychotherapy-oriented childbirth education programme for Chinese first-time childbearing women at 3-month follow up: randomised controlled trial. *International Journal of Nursing Studies*, 49(3), pp.274-81.

Gao L L, Xie W and Yang X ; Chan S W;. (2015). Effects of an interpersonal-psychotherapy-oriented postnatal programme for Chinese first-time mothers: a randomized controlled trial. *International Journal of Nursing Studies*, 52(1), pp.22-9.

Garcia K S, Flynn P and Pierce K J; Caudle M ;. (2010). Repetitive transcranial magnetic stimulation treats postpartum depression. *Brain Stimulation*, 3(1), pp.36-41.

García González, J and Ventura Miranda ; M I ; Requena Mullor ; M ; Parron Carreño ; T ; Alarcón Rodríguez ; R ;. (2018). Effects of prenatal music stimulation on state/trait anxiety in full-term pregnancy and its influence on childbirth: a randomized controlled trial. *Journal of Maternal-Fetal & Neonatal Medicine*, 31(8), pp.1058-1065.

Garnier Salome, De Tychev and Claude ; Lighezzolo Joelle ; Claudon Philippe ; Rebourg-Roesler Christine ; Flach Isabelle ;. (2008). Perinatal depression, prevention strategies, personality, and the importance of therapy acceptance. *Rorschachiana*, 29(2), pp.108-147.

George C. (2019). Effectiveness of a group intervention led by lay health workers in reducing the incidence of postpartum depression in South India. *Asian Journal of Psychiatry*, 47, pp.101864.

Georgiopoulos A M and Bryan T L; Yawn B P; Houston M S; Rummans T A; Therneau T M;. (1999). Population-based screening for postpartum depression. *Obstetrics & Gynecology*, 93(5 Pt 1), pp.653-7.

Georgiopoulos A M, Bryan T L; Wollan P and Yawn B P;. (2001). Routine screening for postpartum depression. *Journal of Family Practice*, 50(2), pp.117-22.

Gerrard J, Holden J M and Elliott S A; McKenzie P ; McKenzie J ; Cox J L;. (1993). A trainer's perspective of an innovative programme teaching health visitors about the detection, treatment and prevention of postnatal depression. *Journal of Advanced Nursing*, 18(11), pp.1825-32.

Gibbon K. (2004). Developments in perinatal mental health assessments. *British Journal of Midwifery*, 12(12), pp.754-760.

Gillis B D and Holley S L; Leming-Lee T S; Parish A L;. (2019). Implementation of a Perinatal Depression Care Bundle in a Nurse-Managed Midwifery Practice. *Nursing for Women's Health*, 23(4), pp.288-298.

Ginsburg G S, Barlow A and Goklish N ; Hastings R ; Baker E V; Mullany B ; Tein J Y; Walkup J ;. (2012). Postpartum Depression Prevention for Reservation-Based American Indians: Results from a Pilot Randomized Controlled Trial. *Child & Youth Care Forum*, 41(3), pp.229-245.

Gjerdingen D, Crow S and McGovern P ; Miner M ; Center B ;. (2009). Postpartum depression screening at well-child visits: validity of a 2-question screen and the PHQ-9. *Annals of Family Medicine*, 7(1), pp.63-70.

Gjerdingen D, Crow S and McGovern P ; Miner M ; Center B ;. (2009). Stepped care treatment of postpartum depression: impact on treatment, health, and work outcomes. *Journal of the American Board of Family Medicine: JABFM*, 22(5), pp.473-82.

Gjerdingen D, McGovern P and Center B ;. (2011). Problems with a diagnostic depression interview in a postpartum depression trial. *Journal of the American Board of Family Medicine: JABFM*, 24(2), pp.187-93.

Gjerdingen D K, McGovern P and Pratt R ; Johnson L ; Crow S ;. (2013). Postpartum doula and peer telephone support for postpartum depression: a pilot randomized controlled trial. *Journal of Primary Care & Community Health*, 4(1), pp.36-43.

Glavin K, Smith L and Sorum R ; Ellefsen B ;. (2010). Redesigned community postpartum care to prevent and treat postpartum depression in women--a one-year follow-up study. *Journal of Clinical Nursing*, 19(21-22), pp.3051-62.

Goodman J H, Guarino A and Chenausky K ; Klein L ; Prager J ; Petersen R ; Forget A ; Freeman M ;. (2014). CALM Pregnancy: results of a pilot study of mindfulness-based cognitive therapy for perinatal anxiety. *Archives of Women's Mental Health*, 17(5), pp.373-87.

Goudarzi Z, Askari M and Seyed-Fatemi N ; Asgari P ; Mehran A ;. (2016). The effect of educational program on stress, anxiety and depression of the mothers of neonates having colostomy. *Journal of Maternal-Fetal & Neonatal Medicine*, 29(23), pp.3902-5.

Gray Lisa A and Price Sarah Kye;. (2014). Partnering for mental health promotion: Implementing evidence based mental health services within a maternal and child home health visiting program. *Clinical Social Work Journal*, 42(1), pp.70-80.

Green S M, Haber E and Frey B N; McCabe R E;. (2015). Cognitive-behavioral group treatment for perinatal anxiety: a pilot study. *Archives of Women's Mental Health*, 18(4), pp.631-8.

Grote Nancy K and Bledsoe Sarah E; Swartz Holly A; Frank Ellen. (2004). Culturally Relevant Psychotherapy for Perinatal Depression in Low-Income Ob/Gyn Patients. *Clinical Social Work Journal*, 32(3), pp.327-347.

Grote Nancy K and Bledsoe Sarah E; Swartz Holly A; Frank Ellen. (2004). Feasibility of Providing Culturally Relevant, Brief Interpersonal Psychotherapy for Antenatal Depression in an Obstetrics Clinic: A Pilot Study. *Research on Social Work Practice*, 14(6), pp.397-407.

Grote N K, Swartz H A; Geibel S L; Zuckoff A and Houck P R; Frank E ;. (2009). A randomized controlled trial of culturally relevant, brief interpersonal psychotherapy for perinatal depression. *Psychiatric services (Washington and D.C.)*, 60(3), pp.313-321.

Grote N K, Katon W J; Lohr M J; Carson K and Curran M ; Galvin E ; Russo J E; Gregory M ;. (2014). Culturally relevant treatment services for perinatal depression in socio-economically disadvantaged women: the design of the MOMCare study. *Contemporary Clinical Trials*, 39(1), pp.34-49.

Gunning M D and Denison F C; Stockley C J; Ho S P; Sandhu H K; Reynolds R M;. (2010). Assessing maternal anxiety in pregnancy with the State-Trait Anxiety Inventory (STAI): Issues of validity, location and participation. *Journal of Reproductive and Infant Psychology*, 28(3), pp.266-273.

Haakstad L A, Torset B and Bo K ;. (2016). What is the effect of regular group exercise on maternal psychological outcomes and common pregnancy complaints? An assessor blinded RCT. *Midwifery*, 32, pp.81-6.

Haga S M, Drozd F and Brendryen H ; Slinning K ;. (2013). Mamma mia: a feasibility study of a web-based intervention to reduce the risk of postpartum depression and enhance subjective well-being. *JMIR Research Protocols*, 2(2), pp.e29.

Haga S M, Drozd F and Lisoy C ; Wentzel-Larsen T ; Slinning K ;. (2019). Mamma Mia - A randomized controlled trial of an internet-based intervention for perinatal depression. *Psychological Medicine*, 49(11), pp.1850-1858.

Hajmohamadi N, Ghalichi F and Bakhtari Aghdam ; F ; Matlabi H ;. (2018). The "Cooperative-Supportive" Intervention for Improving Mental Health Status among Pregnant Women. *Journal of Caring Sciences*, 7(2), pp.101-106.

Hall K and Grundy S . (2014). An analysis of Time 4U, a therapeutic group for women with postnatal depression. *Community Practitioner*, 87(9), pp.25-8.

Hanbury A, Farley K and Thompson C ; Wilson P M; Chambers D ; Holmes H ;. (2013). Immediate versus sustained effects: interrupted time series analysis of a tailored intervention. *Implementation Science*, 8, pp.130.

Hancock H S, Pituch K and Uzark K ; Bhat P ; Fifer C ; Silveira M ; Yu S ; Welch S ; Donohue J ; Lowery R ; Aiyagari R ;. (2018). A randomised trial of early palliative care for maternal stress in infants prenatally diagnosed with single-ventricle heart disease. *Cardiology in the Young*, 28(4), pp.561-570.

Hanna B, Jarman H and Savage S ;. (2004). The clinical application of three screening tools for recognizing post-partum depression. *International Journal of Nursing Practice*, 10(2), pp.72-9.

Hantsoo L, Ward-O'Brien D and Czarkowski K A; Gueorguieva R ; Price L H; Epperson C N;. (2014). A randomized, placebo-controlled, double-blind trial of sertraline for postpartum depression. *Psychopharmacology*, 231(5), pp.939-48.

Hantsoo L, Podcasy J and Sammel M ; Epperson C N; Kim D R;. (2017). Pregnancy and the Acceptability of Computer-Based Versus Traditional Mental Health Treatments. *Journal of Women's Health*, 26(10), pp.1106-1113.

Hantsoo L, Criniti S and Khan A ; Moseley M ; Kincler N ; Faherty L J; Epperson C N; Bennett I M;. (2018). A mobile application for monitoring and management of depressed mood in a vulnerable pregnant population. *Psychiatric Services*, 69(1), pp.104-107.

Harrington A R and Greene-Harrington C C; Harrington Alfred R; Greene-Harrington Claudia C;. (2007). Healthy Start screens for depression among urban pregnant, postpartum and interconceptional women. *Journal of the National Medical Association*, 99(3), pp.226-231.

Harrison P A and Sidebottom A C;. (2008). Systematic prenatal screening for psychosocial risks. *Journal of Health Care for the Poor & Underserved*, 19(1), pp.258-76.

Harrison-Hohner J, Coste S and Dorato V ; Curet L B; McCarron D ; Hatton D ;. (2001). Prenatal calcium supplementation and postpartum depression: An ancillary study to a randomized trial of calcium for prevention of preeclampsia. *Archives of Women's Mental Health*, 3(4), pp.141-146.

Harvey S T and Fisher L J; Green V M;. (2012). Evaluating the clinical efficacy of a primary care-focused, nurse-led, consultation liaison model for perinatal mental health. *International Journal of Mental Health Nursing*, 21(1), pp.75-81.

Harvey Simone T, Bennett Jennifer A; Burmeister Elizabeth and Wyder Marianne ;. (2018). Evaluating a nurse-led community model of service for perinatal mental health. *Collegian*, 25(5), pp.525-531.

Hauck Y, Rock D and Jackiewicz T ; Jablensky A ;. (2008). Healthy babies for mothers with serious mental illness: a case management framework for mental health clinicians. *International Journal of Mental Health Nursing*, 17(6), pp.383-391.

Hayes B A, Muller R and Bradley B S;. (2001). Perinatal depression: a randomized controlled trial of an antenatal education intervention for primiparas. *Birth*, 28(1), pp.28-35.

Heberlein Emily C. (2015). The comparative effectiveness of group prenatal care on women's psychosocial health. *Dissertation Abstracts International: Section B: The Sciences and Engineering*, 75(8-B(E)), pp.No Pagination Specified.

Heberlein E C, Picklesimer A H; Billings D L; Covington-Kolb S and Farber N ; Frongillo E A;. (2016). The comparative effects of group prenatal care on psychosocial outcomes. *Archives of Women's Mental Health*, 19(2), pp.259-69.

Hegarty K, Brown S and Gunn J ; Forster D ; Nagle C ; Grant B ; Lumley J ;. (2007). Women's views and outcomes of an educational intervention designed to enhance psychosocial support for women during pregnancy. *Birth*, 34(2), pp.155-63.

Hepp P, Hagenbeck C and Gilles J ; Wolf O T; Goertz W ; Janni W ; Balan P ; Fleisch M ; Fehm T ; Schaal N K;. (2018). Effects of music intervention during caesarean delivery on anxiety and stress of the mother a controlled, randomised study. *BMC Pregnancy & Childbirth*, 18(1), pp.435.

Herizchi S, Hosseini M B and Ghoreishizadeh M ;. (2017). The impact of Kangaroo-Mother care on postpartum depression in mothers of premature infants. *International Journal of Women's Health and Reproduction Sciences*, 5(4), pp.312-317.

Highet N and Drummond P . (2004). A comparative evaluation of community treatments for post-partum depression: implications for treatment and management practices. *Australian & New Zealand Journal of Psychiatry*, 38(4), pp.212-8.

Highet Nicole, Gamble Jenny and Creedy Debra ;. (2018). Perinatal mental health and psychosocial risk screening in a community maternal and child health setting: Evaluation of a digital platform. *Primary Health Care Research and Development*, , pp.No Pagination Specified.

Hillier Carey A and Slade P. (1989). The impact of antenatal classes on knowledge, anxiety and confidence in primiparous women. *Journal of Reproductive and Infant Psychology*, 7(1), pp.3-13.

Hiscock H, Bayer J and Gold L ; Hampton A ; Ukoumunne O C; Wake M ;. (2007). Improving infant sleep and maternal mental health: a cluster randomised trial. *Archives of Disease in Childhood*, 92(11), pp.952-8.

Ho S M and Heh S S; Jevitt C M; Huang L H; Fu Y Y; Wang L L;. (2009). Effectiveness of a discharge education program in reducing the severity of postpartum depression: a randomized controlled evaluation study. *Patient Education & Counseling*, 77(1), pp.68-71.

Hoffman Melissa. (2019). A screening protocol for perinatal mood and anxiety disorders in expectant mothers. *Dissertation Abstracts International: Section B: The Sciences and Engineering*, 80(5-B(E)), pp.No Pagination Specified.

Holt C, Milgrom J and Gemmill A W;. (2017). Improving help-seeking for postnatal depression and anxiety: a cluster randomised controlled trial of motivational interviewing. *Archives of Women's Mental Health*, 20(6), pp.791-801.

Honey K L, Bennett P and Morgan M ;. (2002). A brief psycho-educational group intervention for postnatal depression. *British Journal of Clinical Psychology*, 41(Pt 4), pp.405-9.

Horowitz J A and Murphy C A; Gregory K E; Wojcik J. (2011). A community-based screening initiative to identify mothers at risk for postpartum depression. *JOGNN - Journal of Obstetric, Gynecologic and Neonatal Nursing*, 40(1), pp.52-61.

Horsch A, Tolsa J F and Gilbert L ; du Chene ; L J ; Muller-Nix C ; Bickle Graz ; M ;. (2016). Improving Maternal Mental Health Following Preterm Birth Using an Expressive Writing Intervention: A Randomized Controlled Trial. *Child Psychiatry & Human Development*, 47(5), pp.780-91.

Hosseini S, Shahtaheri E and Ebrahimi M ; Mahdavi A ; Sepahvandi M A;. (2016). Effectiveness of mindfulness-based cognitive therapy and metacognition therapy on reduction of symptoms of depression, core self-evaluation and emotion regulation difficulties in pregnant women with depression. *Acta Medica Mediterranea*, 32(SpecialIssue5), pp.2033-2037.

Howard M, Battle C L and Pearlstein T ; Rosene-Montella K ;. (2006). A psychiatric mother-baby day hospital for pregnant and postpartum women. *Archives of Women's Mental Health*, 9(4), pp.213-8.

Huang Z J and Wong F Y; Ronzio C R; Yu S M;. (2007). Depressive symptomatology and mental health help-seeking patterns of U.S.- and foreign-born mothers. *Maternal & Child Health Journal*, 11(3), pp.257-67.

Hublely Samuel Henry. (2014). Training obstetrics nurses and behavioral health care providers in behavioral activation for antenatal depression. *Dissertation Abstracts International: Section B: The Sciences and Engineering*, 75(4-B(E)), pp.No Pagination Specified.

Hung K J, Tomlinson M and le Roux ; I M ; Dewing S ; Chopra M ; Tsai A C;. (2014). Community-based prenatal screening for postpartum depression in a South African township. *International Journal of Gynaecology & Obstetrics*, 126(1), pp.74-7.

Ickovics J R, Reed E and Magriples U ; Westdahl C ; Schindler Rising ; S ; Kershaw T S;. (2011). Effects of group prenatal care on psychosocial risk in pregnancy: results from a randomised controlled trial. *Psychology & Health*, 26(2), pp.235-50.

Iheanacho T, Obiefune M and Ezeanolue C O; Ogedegbe G ; Nwanyanwu O C; Ehiri J E; Ohaeri J ; Ezeanolue E E;. (2015). Integrating mental health screening into routine community maternal and child health activity: experience from Prevention of Mother-to-child HIV transmission (PMTCT) trial in Nigeria. *Social Psychiatry & Psychiatric Epidemiology*, 50(3), pp.489-95.

Imran N and Haider I I. (2010). Screening of antenatal depression in Pakistan: Risk factors and effects on obstetric and neonatal outcomes. *Asia-Pacific Psychiatry*, 2(1), pp.26-32.

Imura M, Misao H and Ushijima H ;. (2006). The psychological effects of aromatherapy-massage in healthy postpartum mothers. *Journal of Midwifery & Women's Health*, 51(2), pp.e21-7.

Ingadottir E and Thome M . (2006). Evaluation of a web-based course for community nurses on postpartum emotional distress. *Scandinavian Journal of Caring Sciences*, 20(1), pp.86-92.

Ingram J and Taylor J . (2007). Predictors of postnatal depression: Using an antenatal needs assessment discussion tool. *Journal of Reproductive and Infant Psychology*, 25(3), pp.210-222.

Ip L S and Chau J P. C; Thompson D R; Choi K C;. (2015). An evaluation of a nurse-led Comprehensive Child Development Service in Hong Kong. *Journal of Reproductive and Infant Psychology*, 33(1), pp.88-98.

Jackson Fatimah Zahirah. (2018). Maternal depression in Barbados: Exploring how black women experience, understand, manage and cope with self-reported 'baby blues' and postpartum depression. *Dissertation Abstracts International: Section B: The Sciences and Engineering*, 79(5-B(E)), pp.No Pagination Specified.

Jahdi F, Koohsarian Z and Rasoulia M ; Montazerim A ;. (2016). Evaluating the effectiveness of empowerment-based education on empowerment and anxiety in nulliparous women. *Acta Medica Mediterranea*, 32(SpecialIssue4), pp.1281-1287.



- Jallo N, Cozens R and Smith M W; Simpson R I;. (2013). Effects of a guided imagery intervention on stress in hospitalized pregnant women: a pilot study. *Holistic Nursing Practice*, 27(3), pp.129-39.
- Jardri R, Maron M and Pelta J ; Thomas P ; Codaccioni X ; Goudemand M ; Delion P ;. (2010). Impact of midwives' training on postnatal depression screening in the first week post delivery: a quality improvement report. *Midwifery*, 26(6), pp.622-9.
- Jesse D E, Blanchard A and Bunch S ; Dolbier C ; Hodgson J ; Swanson M S;. (2010). A pilot study to reduce risk for antepartum depression among women in a public health prenatal clinic. *Issues in Mental Health Nursing*, 31(5), pp.355-64.
- Jesse D E, Gaynes B N; Feldhousen E B; Newton E R; Bunch S and Hollon S D;. (2015). Performance of a Culturally Tailored Cognitive-Behavioral Intervention Integrated in a Public Health Setting to Reduce Risk of Antepartum Depression: A Randomized Controlled Trial. *Journal of Midwifery & Women's Health*, 60(5), pp.578-92.
- Jevitt Cecilia, Zapata Lauren and Harrington Monalisa ; Berry Estrellita ;. (2005). Screening for perinatal depression with limited psychiatric resources. *Journal of the American Psychiatric Nurses Association*, 11(6), pp.359-363.
- Jiang L, Wang Z Z and Qiu L R; Wan G B; Lin Y ; Wei Z ;. (2014). Psychological intervention for postpartum depression. *Journal of Huazhong University of Science and Technology. Medical Sciences*, 34(3), pp.437-442.
- Jiao N, Zhu L and Chong Y S; Chan W S; Luo N ; Wang W ; Hu R ; Chan Y H; He H G;. (2019). Web-based versus home-based postnatal psychoeducational interventions for first-time mothers: A randomised controlled trial. *International Journal of Nursing Studies*, 99, pp.103385.
- John H B, Philip R M; Santhanam S and Padankatti S M; Sebastian T ; Balan I ; Rajapandian E ;. (2018). Activity based group therapy reduces maternal anxiety in the Neonatal Intensive Care Unit - a prospective cohort study. *Early Human Development*, 123, pp.17-21.
- Johnson K, Woodward A and Swenson S ; Weis C ; Gunderson M ; Deling M ; Cristiani V ; Lynch B ;. (2017). Parents' adverse childhood experiences and mental health screening using home visiting programs: A pilot study. *Public Health Nursing*, 34(6), pp.522-530.
- Johnson-Agbakwu C E, Allen J and Nizigiyimana J F; Ramirez G ; Hollifield M ;. (2014). Mental health screening among newly arrived refugees seeking routine obstetric and gynecologic care. *Psychological Services*, 11(4), pp.470-476.
- Jomeen J and Martin C R. (2008). The impact of choice of maternity care on psychological health outcomes for women during pregnancy and the postnatal period. *Journal of Evaluation in Clinical Practice*, 14(3), pp.391-8.
- Jones C J and Creedy D K; Gamble J A;. (2012). Australian midwives' awareness and management of antenatal and postpartum depression. *Women & Birth: Journal of the Australian College of Midwives*, 25(1), pp.23-8.
- Jones C C, Jomeen J and Hayter M ;. (2015). A Home-Start peer support scheme for women with low mood following childbirth. *Community Practitioner*, 88(9), pp.41-4.

Kaaya S F, Blander J and Antelman G ; Cyprian F ; Emmons K M; Matsumoto K ; Chopyak E ; Levine M ; Smith Fawzi ; M C ;. (2013). Randomized controlled trial evaluating the effect of an interactive group counseling intervention for HIV-positive women on prenatal depression and disclosure of HIV status. *AIDS Care*, 25(7), pp.854-62.

Kadivar M, Seyedfatemi N and Akbari N ; Haghani H ;. (2015). The effect of narrative writing on maternal stress in neonatal intensive care settings. *Journal of Maternal-Fetal & Neonatal Medicine*, 28(8), pp.938-43.

Kalina Emily Jean. (2016). Standardized postpartum depression screening and treatment. *Dissertation Abstracts International: Section B: The Sciences and Engineering*, 76(9-B(E)), pp.No Pagination Specified.

Kalra H, Reilly N and Austin M P;. (2018). An evaluation of routine antenatal depression screening and psychosocial assessment in a regional private maternity setting in Australia. *Australian & New Zealand Journal of Obstetrics & Gynaecology*, 58(6), pp.629-635.

Kamibeppu K, Furuta M and Yamashita H ; Sugishita K ; Suzumiya H ; Yoshida K ;. (2009). Training health professionals to detect and support mothers at risk of postpartum depression or infant abuse in the community: a cross-sectional and a before and after study. *Bioscience Trends*, 3(1), pp.17-24.

Karamoozian M and Askarizadeh G . (2015). Impact of prenatal cognitive-behavioral stress management intervention on maternal anxiety and depression and newborns' Apgar scores. *Iranian Journal of Neonatology*, 6(2), pp.14-23.

Karatas J C, Matthey S and Barnett B ;. (2009). Antenatal psychosocial assessment: how accurate are we in determining 'low-risk' status? A pilot study. *Archives of Women's Mental Health*, 12(2), pp.97-103.

Kelman Alex R and Evare Benjamin S; Barrera Alinne Z; Muñoz Ricardo F; Gilbert Paul. (2018). A proof-of-concept pilot randomized comparative trial of brief Internet-based compassionate mind training and cognitive-behavioral therapy for perinatal and intending to become pregnant women. *Clinical Psychology & Psychotherapy*, 25(4), pp.608-619.

Kelman A R and Evare B S; Barrera A Z; Munoz R F; Gilbert P. (2018). A proof-of-concept pilot randomized comparative trial of brief Internet-based compassionate mind training and cognitive-behavioral therapy for perinatal and intending to become pregnant women. *Clinical Psychology & Psychotherapy*, 23, pp.23.

Khamseh F, Parandeh A and Hajiamini Z ; Tadrissi S D; Najjar M ;. (2019). Effectiveness of applying problem-solving training on depression in Iranian pregnant women: Randomized clinical trial. *Journal of Education & Health Promotion*, 8, pp.87.

Khan M N, Dherani M and Chiumento A ; Atif N ; Bristow K ; Sikander S ; Rahman A ;. (2017). Evaluating feasibility and acceptability of a local psycho-educational intervention for pregnant women with common mental problems affected by armed conflict in Swat, Pakistan: A parallel randomized controlled feasibility trial. *International Journal of Social Psychiatry*, 63(8), pp.724-735.

Khan Faaiza and Laurent Heidemarie K. (2018). Assessing the impact of mindfulness and life stress on maternal well-being. *Mindfulness*, , pp.No Pagination Specified.

Khan S, Lovell K and Lunat F ; Masood Y ; Shah S ; Tomenson B ; Husain N ;. (2019). Culturally-adapted cognitive behavioural therapy based intervention for maternal depression: a mixed-methods feasibility study. *BMC Women's Health*, 19(1), pp.21.

Khodakarami Batul, Bibalan Faegheh Gotalizadeh and Soltani Farzaneh ; Soltanian Alireza ; Mohagheghi Hussein ;. (2017). Impact of a Counseling Program on Depression, Anxiety, Stress, and Spiritual Intelligence in Pregnant Women. *Journal of Midwifery & Reproductive Health*, 5(2), pp.858-866.

Khojasteh F, Rezaee N and Safarzadeh A ; Sahlabadi R ; Shahrakipoor M ;. (2016). Comparison of the effects of massage therapy and guided imagery on anxiety of nulliparous women during pregnancy. *Der Pharmacia Lettre*, 8(19), pp.1-7.

Kildea S, Simcock G and Liu A ; Elgbeili G ; Laplante D P; Kahler A ; Austin M P; Tracy S ; Kruske S ; Tracy M ; O'Hara M W; King S ;. (2018). Continuity of midwifery carer moderates the effects of prenatal maternal stress on postnatal maternal wellbeing: the Queensland flood study. *Archives of Women's Mental Health*, 21(2), pp.203-214.

Kim H, Bracha Y and Tipnis A ;. (2007). Automated depression screening in disadvantaged pregnant women in an urban obstetric clinic. *Archives of Women's Mental Health*, 10(4), pp.163-9.

Kim J J, La Porte and L M ; Corcoran M ; Magasi S ; Batza J ; Silver R K;. (2010). Barriers to mental health treatment among obstetric patients at risk for depression. *American Journal of Obstetrics & Gynecology*, 202(3), pp.312.e1-5.

Kim D R, Sockol L and Barber J P; Moseley M ; Lamprou L ; Rickels K ; O'Reardon J P; Epperson C N;. (2011). A survey of patient acceptability of repetitive transcranial magnetic stimulation (TMS) during pregnancy. *Journal of Affective Disorders*, 129(1-3), pp.385-90.

Kim H G, Geppert J and Quan T ; Bracha Y ; Lupo V ; Cutts D B;. (2012). Screening for postpartum depression among low-income mothers using an interactive voice response system. *Maternal & Child Health Journal*, 16(4), pp.921-8.

Kim D R, Hantsoo L and Thase M E; Sammel M ; Epperson C N;. (2014). Computer-assisted cognitive behavioral therapy for pregnant women with major depressive disorder. *Journal of Women's Health*, 23(10), pp.842-8.

Kimmel M C, Cluxton-Keller F and Frosch E ; Carter T ; Solomon B S;. (2017). Maternal Experiences in a Parenting Group Delivered in an Urban General Pediatric Clinic. *Clinical Pediatrics*, 56(1), pp.13-19.

Kingston D E, Biringner A and Toosi A ; Heaman M I; Lasiuk G C; McDonald S W; Kingston J ; Sword W ; Jarema K ; Austin M P;. (2015). Disclosure during prenatal mental health screening. *Journal of Affective Disorders*, 186, pp.90-4.

Kingston D, Austin M P and Heaman M ; McDonald S ; Lasiuk G ; Sword W ; Giallo R ; Hegadoren K ; Vermeyden L ; van Zanten ; S V ; Kingston J ; Jarema K ; Biringner A ;. (2015). Barriers and facilitators of mental health screening in pregnancy. *Journal of Affective Disorders*, 186, pp.350-7.

Kingston D, Biringner A and Veldhuyzen van Zanten; S ; Giallo R ; McDonald S ; MacQueen G ; Vermeyden L ; Austin M P;. (2017). Pregnant Women's Perceptions of the Risks and

Benefits of Disclosure During Web-Based Mental Health E-Screening Versus Paper-Based Screening: Randomized Controlled Trial. *JMIR Mental Health*, 4(4), pp.e42.

Kingston D, Austin M P and Veldhuyzen van Zanten; S ; Harvalik P ; Giallo R ; McDonald S D; MacQueen G ; Vermeyden L ; Lasiuk G ; Sword W ; Biringer A ;. (2017). Pregnant Women's Views on the Feasibility and Acceptability of Web-Based Mental Health E-Screening Versus Paper-Based Screening: A Randomized Controlled Trial. *Journal of Medical Internet Research*, 19(4), pp.e88.

Kleiber B V, Felder J N; Ashby B and Scott S ; Dean J ; Dimidjian S ;. (2017). Treating Depression Among Adolescent Perinatal Women With a Dialectical Behavior Therapy-Informed Skills Group. *Cognitive and Behavioral Practice*, 24(4), pp.416-427.

Ko Y and Yang C . (2008). Effects of postpartum exercise program on fatigue and depression during 'doing-the-month' period. *Journal of Nursing Research (Taiwan Nurses Association)*, 16(3), pp.177-185.

Ko Yi-Li, Yang Chi-Li and Fang Chin-Lung ; Lee Mei-Ying ; Lin Pi-Chu ;. (2013). Community-based postpartum exercise program. *Journal of Clinical Nursing (John Wiley & Sons and Inc.)*, 22(15-16), pp.2122-2131.

Kohlhoff J, Hickinbotham R and Knox C ; Roach V ; Barnett Am ; B ;. (2016). Antenatal psychosocial assessment and depression screening in a private hospital. *Australian & New Zealand Journal of Obstetrics & Gynaecology*, 56(2), pp.173-8.

Kopelman R C, Moel J and Mertens C ; Stuart S ; Arndt S ; O'Hara M W;. (2008). Barriers to care for antenatal depression. *Psychiatric Services*, 59(4), pp.429-32.

Kozinszky Z, Dudas R B and Devosa I ; Csator dai S ; Toth E ; Szabo D ; Sikovanyecz J ; Barabas K ; Pal A ;. (2012). Can a brief antepartum preventive group intervention help reduce postpartum depressive symptomatology?. *Psychotherapy & Psychosomatics*, 81(2), pp.98-107.

Kralj Holly T. (2016). Creating pathways of care: A multidisciplinary approach to increasing screening and treatment of postpartum depression within a rural health clinic. *Dissertation Abstracts International: Section B: The Sciences and Engineering*, 76(8-B(E)), pp.No Pagination Specified.

Krusche A, Dymond M and Murphy S E; Crane C ;. (2018). Mindfulness for pregnancy: A randomised controlled study of online mindfulness during pregnancy. *Midwifery*, 65, pp.51-57.

Kurzweil S. (2012). Psychodynamic therapy for depression in women with infants and young children. *American Journal of Psychotherapy*, 66(2), pp.181-99.

La Porte , L M and Kim J J; Adams M G; Zagorsky B M; Gibbons R ; Silver R K;. (2019). Feasibility of perinatal mood screening and text messaging on patients' personal smartphones. *Archives of Women's Mental Health*, 15, pp.15.

Laios L, Steele A and Judd F ;. (2010). What are we screening for? Development of a psychiatric referral tool for use with adolescent pregnant women. *Australasian Psychiatry*, 18(3), pp.256-60.

Langer A, Campero L and Garcia C ; Reynoso S ;. (1998). Effects of psychosocial support during labour and childbirth on breastfeeding, medical interventions, and mothers' wellbeing in a Mexican public hospital: a randomised clinical trial. *British Journal of Obstetrics & Gynaecology*, 105(10), pp.1056-63.

Lara M A, Navarro C and Navarrete L ;. (2010). Outcome results of a psycho-educational intervention in pregnancy to prevent PPD: a randomized control trial. *Journal of Affective Disorders*, 122(1-2), pp.109-17.

Lara Ma, Navarro Claudia and Navarrete Laura ; Le Huynh-Nhu ;. (2010). Retention rates and potential predictors in a longitudinal randomized control trial to prevent postpartum depression. *Salud Mental*, 33(5), pp.429-436.

Lara Ma, Navarrete Laura and Nieto Lourdes ; Berenzon Shoshana ;. (2014). Acceptability and barriers to treatment for perinatal depression. An exploratory study in Mexican women. *Salud Mental*, 37(4), pp.293-301.

LaRocco-Cockburn A, Reed S D and Melville J ; Croicu C ; Russo J E; Inspektor M ; Edmondson E ; Katon W ;. (2013). Improving depression treatment for women: integrating a collaborative care depression intervention into OB-GYN care. *Contemporary Clinical Trials*, 36(2), pp.362-70.

Lawson A, Dalfen A and Murphy K E; Milligan N ; Lancee W ;. (2019). Use of Text Messaging for Postpartum Depression Screening and Information Provision. *Psychiatric Services*, 70(5), pp.389-395.

Le H N and Perry D F; Sheng X. (2009). Using the internet to screen for postpartum depression. *Maternal & Child Health Journal*, 13(2), pp.213-21.

Le H N, Zmuda J and Perry D F; Munoz R F;. (2010). Transforming an evidence-based intervention to prevent perinatal depression for low-income Latina immigrants. *American Journal of Orthopsychiatry*, 80(1), pp.34-45.

Le H N, Perry D F; Genovez M and Cardeli E ;. (2013). In their own voices: Latinas' experiences with a randomized controlled trial. *Qualitative Health Research*, 23(6), pp.834-46.

Le H N, Rodriguez-Munoz M F; Soto-Balbuena C and Olivares Crespo ; M E ; Izquierdo Mendez ; N ; Marcos-Najera R ;. (2019). Preventing perinatal depression in Spain: a pilot evaluation of Mamas y Bebés. *Journal of Reproductive & Infant Psychology*, , pp.1-14.

Leigh B and Milgrom J . (2007). Acceptability of antenatal screening for depression in routine antenatal care. *Australian Journal of Advanced Nursing*, 24(3), pp.14-8.

Leis J A, Mendelson T and Perry D F; Tandon S D;. (2011). Perceptions of mental health services among low-income, perinatal African-American women. *Womens Health Issues*, 21(4), pp.314-9.

Lenze S N, Rodgers J and Luby J ;. (2015). A pilot, exploratory report on dyadic interpersonal psychotherapy for perinatal depression. *Archives of Women's Mental Health*, 18(3), pp.485-91.

Lenze S N and Potts M A;. (2017). Brief Interpersonal Psychotherapy for depression during pregnancy in a low-income population: A randomized controlled trial. *Journal of Affective Disorders*, 210, pp.151-157.

Letourneau N, Duffett-Leger L and Stewart M ; Hegadoren K ; Dennis C L; Rinaldi C M; Stoppard J ;. (2007). Canadian mothers' perceived support needs during postpartum depression. *JOGNN - Journal of Obstetric, Gynecologic and Neonatal Nursing*, 36(5), pp.441-9.

Leung Sharron Shuk Kam. (2002). Postpartum depression: Perceived social support and stress among Hong Kong Chinese women. *Dissertation Abstracts International: Section B: The Sciences and Engineering*, 62(10-B), pp.4469.

Leung S S and Lee A M; Chiang V C; Lam S K; Kuen Y W; Wong D F;. (2013). Culturally sensitive, preventive antenatal group cognitive-behavioural therapy for Chinese women with depression. *International Journal of Nursing Practice*, 19 Suppl 1, pp.28-37.

Lewis N L. (2019). Developing a Hospital-Based Postpartum Depression Education Intervention for Perinatal Nurses. *Journal for Nurses in Professional Development*, 14, pp.14.

Lia-Hoagberg Betty, Rode Peter and Skovholt Catherine J; Oberg Charles N; Berg Cynthia ; Mullett Sara ; Choi Thomas ;. (1990). Barriers and motivators to prenatal care among low-income women. *Social Science & Medicine*, 30(4), pp.487-495.

Liberto Terri L. (2011). *Attitudes Toward Help Seeking Among Depressed and Non-depressed Postpartum Women*. . West Virginia University.

Liu Y H and Chang M Y; Chen C H;. (2010). Effects of music therapy on labour pain and anxiety in Taiwanese first-time mothers. *Journal of Clinical Nursing*, 19(7-8), pp.1065-72.

Liu Y Q and Maloni J A; Petrini M A;. (2014). Effect of postpartum practices of doing the month on Chinese women's physical and psychological health. *Biological Research for Nursing*, 16(1), pp.55-63.

Logsdon M C, Cross R and Williams B ; Simpson T ;. (2004). Prediction of postpartum social support and symptoms of depression in pregnant adolescents: a pilot study. *Journal of School Nursing*, 20(1), pp.36-42.

Logsdon M C, Foltz M P; Stein B and Usui W ; Josephson A ;. (2010). Adapting and testing telephone-based depression care management intervention for adolescent mothers. *Archives of Women's Mental Health*, 13(4), pp.307-17.

Logsdon M C, Barone M and Lynch T ; Robertson A ; Myers J ; Morrison D ; York S ; Gregg J ;. (2013). Testing of a prototype Web based intervention for adolescent mothers on postpartum depression. *Applied Nursing Research*, 26(3), pp.143-5.

Logsdon M C, Davis D W; Stikes R and Ratterman R ; Ryan L ; Myers J ;. (2015). Acceptability and initial efficacy of education for teen mothers. *MCN and American Journal of Maternal Child Nursing*, 40(3), pp.186-92.

Logsdon M C, Vogt K and Davis D W; Myers J ; Hogan F ; Eckert D ; Masterson K ;. (2018). Screening for Postpartum Depression by Hospital-Based Perinatal Nurses. *MCN and American Journal of Maternal Child Nursing*, 43(6), pp.324-329.

Lopez A A and Skelly J M; Higgins S T;. (2015). Financial incentives for smoking cessation among depression-prone pregnant and newly postpartum women: effects on smoking abstinence and depression ratings. *Nicotine & Tobacco Research*, 17(4), pp.455-62.

Loudon H, Nentin F and Silverman M E;. (2016). Using clinical decision support as a means of implementing a universal postpartum depression screening program. *Archives of Women's Mental Health*, 19(3), pp.501-5.

Loughnan S A, Sie A and Hobbs M J; Joubert A E; Smith J ; Haskelberg H ; Mahoney A E. J; Kladnitski N ; Holt C J; Milgrom J ; Austin M P; Andrews G ; Newby J M;. (2019). A randomized controlled trial of 'MUMentum Pregnancy': Internet-delivered cognitive behavioral therapy program for antenatal anxiety and depression. *Journal of Affective Disorders*, 243, pp.381-390.

Lovero K L, Lammie S L; van Zyl and A ; Paul S N; Ngwepe P ; Mootz J J; Carlson C ; Sweetland A C; Shelton R C; Wainberg M L; Medina-Marino A ;. (2019). Mixed-methods evaluation of mental healthcare integration into tuberculosis and maternal-child healthcare services of four South African districts. *BMC Health Services Research*, 19(1), pp.83.

Ludwick Susan Danielle. (2017). Postpartum Depression: Implementing an Evidence-Based Social Support Network in North Carolina. *Postpartum Depression: Implementing an Evidence-Based Social Support Network in North Carolina*, , pp.1-1.

MacArthur C, Winter H R and Bick D E; Lilford R J; Lancashire R J; Knowles H ; Braunholtz D A; Henderson C ; Belfield C ; Gee H ;. (2003). Redesigning postnatal care: a randomised controlled trial of protocol-based midwifery-led care focused on individual women's physical and psychological health needs. *Health Technology Assessment (Winchester and England)*, 7(37), pp.1-98.

Madden D, Sliney A and O'Friel A ; McMackin B ; O'Callaghan B ; Casey K ; Courtney L ; Fleming V ; Brady V ;. (2018). Using action research to develop midwives' skills to support women with perinatal mental health needs. *Journal of Clinical Nursing*, 27(3-4), pp.561-571.

Madhavanprabhakaran G K and D'Souza M S; Nairy K S;. (2015). Prevalence of pregnancy anxiety and associated factors. *International Journal of Africa Nursing Sciences*, 3, pp.1-7.

Magalhaes P V, Pinheiro R T; Faria A D; Osorio C M; da Silva and R A ; Botella L ;. (2007). Impact of defense style on brief psychotherapy of postpartum depression. *Journal of Nervous & Mental Disease*, 195(10), pp.870-3.

Mahato P K, van Teijlingen and E ; Simkhada P ; Angell C ; Ireland J ; team Thet ;. (2018). Qualitative evaluation of mental health training of auxiliary nurse midwives in rural Nepal. *Nurse Education Today*, 66, pp.44-50.

Mak J K. L, Lee A H; Pham N M; Tang L and Pan X F; Xu Z P; Binns C W; Sun X ;. (2019). Physical activity during early pregnancy and antenatal depression: A prospective cohort study. *Mental Health and Physical Activity*, 16, pp.54-59.

- Malis Françoise Roy, Meyer Thorsten and Gross Mechthild M; Roy Malis ; Françoise ;. (2017). Effects of an antenatal mindfulness-based childbirth and parenting programme on the postpartum experiences of mothers: a qualitative interview study. *BMC Pregnancy & Childbirth*, 17, pp.1-11.
- Maloni J A, Przeworski A and Damato E G;. (2013). Web recruitment and internet use and preferences reported by women with postpartum depression after pregnancy complications. *Archives of Psychiatric Nursing*, 27(2), pp.90-5.
- Malqvist M, Clarke K and Matsebula T ; Bergman M ; Tomlinson M ;. (2016). Screening for Antepartum Depression Through Community Health Outreach in Swaziland. *Journal of Community Health*, 41(5), pp.946-52.
- Mane Kusum, S and Suryakantha A H;. (2013). A study on mental health status among pregnant women and the social factors influencing. *Indian Journal of Public Health Research and Development*, 4(4), pp.79-83.
- Mann Rachel, Adamson Joy and Gilbody Simon ;. (2015). The acceptability of case-finding questions to identify perinatal depression. *British Journal of Midwifery*, 23(9), pp.630-638.
- Mao Hong-Jing, Li He-Jiang and Chiu Helen ; Chan Wai-Chi ; Chen Shu-Ling ;. (2012). Effectiveness of Antenatal Emotional Self-Management Training Program in Prevention of Postnatal Depression in Chinese Women. *Perspectives in Psychiatric Care*, 48(4), pp.218-224.
- Marcus S M and Flynn H A; Blow F C; Barry K L;. (2003). Depressive symptoms among pregnant women screened in obstetrics settings. *Journal of Women's Health*, 12(4), pp.373-80.
- Marrs J. (2013). Evaluation of the impact of a CBT-based group on maternal postnatal mental health difficulties. *Journal of Reproductive and Infant Psychology*, 31(1), pp.72-80.
- Martin P R, Arria A M; Fischer G and Kaltenbach K ; Heil S H; Stine S M; Coyle M G; Selby P ; Jones H E;. (2009). Psychopharmacologic management of opioid-dependent women during pregnancy. *American Journal on Addictions*, 18(2), pp.148-56.
- Martinez P, Vohringer P A and Rojas G ;. (2016). Barriers to access to treatment for mothers with postpartum depression in primary health care centers: a predictive model. *Revista Latino-Americana de Enfermagem*, 24, pp.e2675.
- Mathew R M, Anju Philip and T ; Sreejamol M G;. (2017). Perceived postpartum stress and coping strategies among postnatal mothers at aims, Kochi. *Asian Journal of Pharmaceutical and Clinical Research*, 10(12), pp.116-119.
- Matsumoto K, Sato K and Hamatani S ; Shirayama Y ; Shimizu E ;. (2019). Cognitive behavioral therapy for postpartum panic disorder: a case series. *BMC psychology*, 7(1), pp.53.
- Matthey S, Della Vedova and A M ;. (2018). A comparison of two measures to screen for emotional health difficulties during pregnancy. *Journal of Reproductive & Infant Psychology*, 36(5), pp.463-475.



- Mazzoni S, Hill P and Briggs A ; Barbier K ; Cahill A ; Macones G ; Colditz G ; Tuuli M ; Carter E ;. (2018). The effect of group prenatal care for women with diabetes on social support and depressive symptoms: a pilot randomized trial. *Journal of Maternal-Fetal & Neonatal Medicine*, , pp.1-6.
- McCabe-Beane J E and Stasik-O'Brien S M; Segre L S;. (2018). Anxiety Screening During Assessment of Emotional Distress in Mothers of Hospitalized Newborns. *JOGNN - Journal of Obstetric, Gynecologic and & Neonatal Nursing*, 47(1), pp.105-113.
- McCarter Deborah E, Demidenko Eugene and Hegel Mark T;. (2018). Measuring outcomes of digital technology-assisted nursing postpartum: A randomized controlled trial. *Journal of Advanced Nursing*, 74(9), pp.2207-2217.
- McCarter-Spaulding D and Shea S . (2016). Effectiveness of Discharge Education on Postpartum Depression. *MCN and American Journal of Maternal Child Nursing*, 41(3), pp.168-72.
- McCarthy M and McMahon C . (2008). Acceptance and experience of treatment for postnatal depression in a community mental health setting. *Health Care for Women International*, 29(6), pp.618-37.
- McFarlane E, Burrell L and Duggan A ; Tandon D ;. (2017). Outcomes of a Randomized Trial of a Cognitive Behavioral Enhancement to Address Maternal Distress in Home Visited Mothers. *Maternal & Child Health Journal*, 21(3), pp.475-484.
- McGregor M, Coghlan M and Dennis C L;. (2014). The effect of physician-based cognitive behavioural therapy among pregnant women with depressive symptomatology: a pilot quasi-experimental trial. *Early intervention in psychiatry*, 8(4), pp.348-57.
- McKean M, Caughey A B and Yuracko McKean ; M A ; Cabana M D; Flaherman V J;. (2018). Postpartum Depression: When Should Health Care Providers Identify Those at Risk?. *Clinical Pediatrics*, 57(6), pp.689-693.
- McKee M, Zayas Luis H and Fletcher Jason ; Boyd Rhonda C; Nam Sung Hee;. (2006). Results of an Intervention to Reduce Perinatal Depression Among Low-Income Minority Women in Community Primary Care. *Journal of Social Service Research*, 32(4), pp.63-81.
- McKellar Lois, Steen Mary and Lorensuhewa Nirtal ;. (2017). Capture my mood: a feasibility study to develop a visual scale for women to self-monitor their mental wellbeing following birth. *Evidence Based Midwifery*, 15(2), pp.54-59.
- McLeish Jenny and Redshaw Maggie . (2017). Mothers' accounts of the impact on emotional wellbeing of organised peer support in pregnancy and early parenthood: a qualitative study. *BMC Pregnancy & Childbirth*, 17, pp.1-14.
- McLeish Jenny and Redshaw Maggie . (2019). "Being the best person that they can be and the best mum": a qualitative study of community volunteer doula support for disadvantaged mothers before and after birth in England. *BMC Pregnancy & Childbirth*, 19(1), pp.N.PAG-N.PAG.
- Mehta D, Quast C and Fasching P A; Seifert A ; Voigt F ; Beckmann M W; Faschingbauer F ; Burger P ; Ekici A B; Kornhuber J ; Binder E B; Goecke T W;. (2012). The 5-HTTLPR

polymorphism modulates the influence on environmental stressors on peripartum depression symptoms. *Journal of Affective Disorders*, 136(3), pp.1192-7.

Meijssen Dominique, Wolf Marie-Jeanne and Koldewijn Karen ; van Baar ; Anneloes ; Kok Joke ;. (2011). Maternal psychological distress in the first two years after very preterm birth and early intervention. *Early Child Development and Care*, 181(1), pp.1-11.

Meine K. (2018). Pregnancy unshackled: Increasing equity through implementation of perinatal depression screening, shared decision making, and treatment for incarcerated women. *Nursing Forum*, 53(4), pp.437-447.

Melnyk B M and Crean H F; Feinstein N F; Fairbanks E. (2008). Maternal anxiety and depression after a premature infant's discharge from the neonatal intensive care unit: explanatory effects of the creating opportunities for parent empowerment program. *Nursing Research*, 57(6), pp.383-94.

Melnyk B M and Feinstein N F;. (2009). Reducing hospital expenditures with the COPE (Creating Opportunities for Parent Empowerment) program for parents and premature infants: an analysis of direct healthcare neonatal intensive care unit costs and savings. *Nursing Administration Quarterly*, 33(1), pp.32-7.

Mendelson T, Leis J A and Perry D F; Stuart E A; Tandon S D;. (2013). Impact of a preventive intervention for perinatal depression on mood regulation, social support, and coping. *Archives of Women's Mental Health*, 16(3), pp.211-8.

Mendelson T, McAfee C and Damian A J; Brar A ; Donohue P ; Sibinga E ;. (2018). A mindfulness intervention to reduce maternal distress in neonatal intensive care: a mixed methods pilot study. *Archives of Women's Mental Health*, 21(6), pp.791-799.

Mgonja S and Schoening A . (2017). Postpartum Depression Screening at Well-Child Appointments: A Quality Improvement Project. *Journal of Pediatric Health Care*, 31(2), pp.178-183.

Mianaei S J, Karahroudy F A; Rassouli M and Tafreshi M Z;. (2014). The effect of Creating Opportunities for Parent Empowerment program on maternal stress, anxiety, and participation in NICU wards in Iran. *Iranian Journal of Nursing and Midwifery Research*, 19(1), pp.94-100.

Miklowitz D J, Semple R J; Hauser M and Elkun D ; Weintraub M J; Dimidjian S ;. (2015). Mindfulness-Based Cognitive Therapy for Perinatal Women with Depression or Bipolar Spectrum Disorder. *Cognitive Therapy and Research*, 39(5), pp.590-600.

Milgrom J, Schembri C and Ericksen J ; Ross J ; Gemmill A W;. (2011). Towards parenthood: an antenatal intervention to reduce depression, anxiety and parenting difficulties. *Journal of Affective Disorders*, 130(3), pp.385-94.

Milgrom J, Holt C and Holt C J; Ross J ; Ericksen J ; Gemmill A W;. (2015). Feasibility study and pilot randomised trial of an antenatal depression treatment with infant follow-up. *Archives of Women's Mental Health*, 18(5), pp.717-30.

Milgrom J, Danaher B G and Gemmill A W; Holt C ; Holt C J; Seeley J R; Tyler M S; Ross J ; Ericksen J ;. (2016). Internet Cognitive Behavioral Therapy for Women With Postnatal

Depression: A Randomized Controlled Trial of MumMoodBooster. *Journal of Medical Internet Research*, 18(3), pp.e54.

Miller L S and Boyd B J; Chernov A J;. (2004). Improving the identification and treatment of postpartum depression in a managed care organization. *Journal of Clinical Outcomes Management*, 11(3), pp.157-161.

Miller L, Gur M and Shanok A ; Weissman M ;. (2008). Interpersonal psychotherapy with pregnant adolescents: two pilot studies. *Journal of Child Psychology & Psychiatry & Allied Disciplines*, 49(7), pp.733-42.

Miller L, Shade M and Vasireddy V ;. (2009). Beyond screening: assessment of perinatal depression in a perinatal care setting. *Archives of Women's Mental Health*, 12(5), pp.329-34.

Miller L J, McGlynn A and Suberlak K ; Rubin L H; Miller M ; Pirec V ;. (2012). Now what? Effects of on-site assessment on treatment entry after perinatal depression screening. *Journal of Women's Health*, 21(10), pp.1046-52.

Misri S, Kostaras X and Fox D ; Kostaras D ;. (2000). The impact of partner support in the treatment of postpartum depression. *Canadian Journal of Psychiatry - Revue Canadienne de Psychiatrie*, 45(6), pp.554-8.

Misri S, Reebye P and Millis L ; Shah S ;. (2006). The impact of treatment intervention on parenting stress in postpartum depressed mothers: a prospective study. *American Journal of Orthopsychiatry*, 76(1), pp.115-9.

Misri S, Eng A B and Abizadeh J ; Blackwell E ; Spidel A ; Oberlander T F;. (2013). Factors impacting decisions to decline or adhere to antidepressant medication in perinatal women with mood and anxiety disorders. *Depression & Anxiety*, 30(11), pp.1129-36.

Misund A R, Nerdrum P and Braten S ; Pripp A H; Diseth T H;. (2013). Long-term risk of mental health problems in women experiencing preterm birth: a longitudinal study of 29 mothers. *Annals of General Psychiatry*, 12(1), pp.33.

Misund A R, Nerdrum P and Diseth T H;. (2014). Mental health in women experiencing preterm birth. *BMC Pregnancy & Childbirth*, 14, pp.263.

Moayedoddin A, Moser D and Nanzer N ;. (2013). The impact of brief psychotherapy centred on parenthood on the anxio-depressive symptoms of mothers during the perinatal period. *Swiss Medical Weekly*, 143, pp.w13769.

Mojrian Marzieh, Alidoosti Kobra and Tirgari Batool ; Mehdizadeh Ali ; Jahani Yones ;. (2018). The Effect of Supportive Counseling on the Symptoms of Acute Stress Disorder Following Emergency Cesarean Section. *Journal of Midwifery & Reproductive Health*, 6(2), pp.1208-1214.

Morrell C J, Ricketts T and Tudor K ; Williams C ; Curran J ; Barkham M ;. (2011). Training health visitors in cognitive behavioural and person-centred approaches for depression in postnatal women as part of a cluster randomised trial and economic evaluation in primary care: the PoNDER trial. *Primary Health Care Research & Development*, 12(1), pp.11-20.

Morris J B. (1987). Group psychotherapy for prolonged postnatal depression. *British Journal of Medical Psychology*, 60(Pt 3), pp.279-81.

- Morton A and Forsey P . (2013). My Time, My Space (an arts-based group for women with postnatal depression): a project report. *Community Practitioner*, 86(5), pp.31-4.
- Mouradian L E and DeGrace B W; Thompson D M;. (2013). Art-based occupation group reduces parent anxiety in the neonatal intensive care unit: a mixed-methods study. *American Journal of Occupational Therapy*, 67(6), pp.692-700.
- Mugweni E, Goodliffe S and Adams C ; Walker M ; Kendall S ;. (2019). "I'll look after the kids while you go and have a shower": an evaluation of a service to address mild to moderate maternal perinatal mental health problems. *Journal of Mental Health*, 28(3), pp.324-330.
- Mundorf C, Shankar A and Moran T ; Heller S ; Hassan A ; Harville E ; Lichtveld M ;. (2018). Reducing the Risk of Postpartum Depression in a Low-Income Community Through a Community Health Worker Intervention. *Maternal & Child Health Journal*, 22(4), pp.520-528.
- Munoz R F, Le H N; Ippen C G; Diaz M A; Urizar Jr and G G ; Soto J ; Mendelson T ; Delucchi K ; Lieberman A F;. (2007). Prevention of Postpartum Depression in Low-Income Women: Development of the Mamas y Bebés/Mothers and Babies Course. *Cognitive and Behavioral Practice*, 14(1), pp.70-83.
- Murphey C, Carter P and Price L R; Champion J D; Nichols F ;. (2017). Psychological Distress in Healthy Low-Risk First-Time Mothers during the Postpartum Period: An Exploratory Study. *Nursing Research and Practice*, 2017, pp.8415083.
- Muzik M, Hamilton S E and Lisa Rosenblum ; K ; Waxler E ; Hadi Z ;. (2012). Mindfulness yoga during pregnancy for psychiatrically at-risk women: preliminary results from a pilot feasibility study. *Complementary Therapies in Clinical Practice*, 18(4), pp.235-40.
- Muzik Maria, Rosenblum Katherine and Alfafara Emily ; Schuster Melisa ; Miller Nicole ; Waddell Rachel ; Kohler Emily ;. (2015). Mom Power: preliminary outcomes of a group intervention to improve mental health and parenting among high-risk mothers. *Archives of Women's Mental Health*, 18(3), pp.507-521.
- Myors K A, Schmied V and Johnson M ; Cleary M ;. (2014). 'My special time': Australian women's experiences of accessing a specialist perinatal and infant mental health service. *Health & Social Care in the Community*, 22(3), pp.268-77.
- Myors K A, Cleary M and Johnson M ; Schmied V ;. (2018). 'Modelling a Secure-Base' for Women with Complex Needs: Attachment-Based Interventions Used by Perinatal and Infant Mental Health Clinicians. *Issues in Mental Health Nursing*, 39(3), pp.226-232.
- Nagle U and Farrelly M . (2018). Women's views and experiences of having their mental health needs considered in the perinatal period. *Midwifery*, 66, pp.79-87.
- Nasiri S, Kordi M and Gharavi M M;. (2015). A comparative study of the effects of problem-solving skills training and relaxation on the score of self-esteem in women with postpartum depression. *Iranian Journal of Nursing and Midwifery Research*, 20(1), pp.105-12.
- Navaie-Waliser M, Martin S L and Tessaro I ; Campbell M K; Cross A W;. (2000). Social support and psychological functioning among high-risk mothers: the impact of the Baby Love Maternal Outreach Worker Program. *Public Health Nursing*, 17(4), pp.280-291.

Navidian A, Sarasiyabi A S and Koochakzai M ;. (2017). The effect of home-based supportive-educational counseling on primigravidas' postpartum stress. *International Journal of Women's Health and Reproduction Sciences*, 5(2), pp.112-118.

Navodani T, Gartland D and Brown S J; Riggs E ; Yelland J ;. (2019). Common maternal health problems among Australian-born and migrant women: A prospective cohort study. *PLoS ONE [Electronic Resource]*, 14(2), pp.e0211685.

Naysmith C, Wells M and Newson S ; Webb J ;. (2015). Development and outcomes of a therapeutic group for women with postnatal depression. *Community Practitioner*, 88(3), pp.35-8.

Ngai F W and Chan S W; Ip W Y;. (2009). The effects of a childbirth psychoeducation program on learned resourcefulness, maternal role competence and perinatal depression: a quasi-experiment. *International Journal of Nursing Studies*, 46(10), pp.1298-306.

Ngai Fei-Wan and Chan Sally Wai-Chi. (2011). Psychosocial factors and maternal wellbeing: An exploratory path analysis. *International Journal of Nursing Studies*, 48(6), pp.725-731.

Ngai F W and Chan S W;. (2012). Learned resourcefulness, social support, and perinatal depression in chinese mothers. *Nursing Research*, 61(2), pp.78-85.

Nguyen P H, DiGirolamo A M; Gonzalez-Casanova I and Pham H ; Hao W ; Nguyen H ; Truong T V; Nguyen S ; Harding K B; Reinhart G A; Martorell R ; Ramakrishnan U ;. (2017). Impact of preconceptional micronutrient supplementation on maternal mental health during pregnancy and postpartum: results from a randomized controlled trial in Vietnam. *BMC Women's Health*, 17(1), pp.44.

Nieminen K, Berg I and Frankenstein K ; Viita L ; Larsson K ; Persson U ; Spanberger L ; Wretman A ; Silfvernagel K ; Andersson G ; Wijma K ;. (2016). Internet-provided cognitive behaviour therapy of posttraumatic stress symptoms following childbirth-a randomized controlled trial. *Cognitive Behaviour Therapy*, 45(4), pp.287-306.

Ningrum T P and Ginanjar A S; Sari Y R;. (2018). Stress management intervention for women with multiple roles: Case study of working women in post-partum period. In: , ed., *Diversity in unity: Perspectives from psychology and behavioral sciences*. New York, NY: Routledge/Taylor & Francis Group; US, pp.27-33.

Nishi D, Su K P and Usuda K ; Chang J P; Chiang Y J; Guu T W; Hamazaki K ; Nakaya N ; Sone T ; Hashimoto K ; Hamazaki T ; Matsuoka Y J;. (2019). Differences between Japan and Taiwan in the treatment of pregnant women with depressive symptoms by omega-3 fatty acids: An open-label pilot study. *Nutritional Neuroscience*, 22(1), pp.63-71.

Njoku Francisca. (2017). Translating Evidence of Skin-to-Skin and Rooming-in to Practice. *Translating Evidence of Skin-to-Skin & Rooming-in to Practice*, , pp.1-1.

Noonan M, Galvin R and Jomeen J ; Doody O ;. (2019). Public health nurses' perinatal mental health training needs: A cross sectional survey. *Journal of Advanced Nursing*, 02, pp.02.

Norouzi F, Keshavarz M and SeyedFatemi N ; Montazeri A ;. (2013). The impact of kangaroo care and music on maternal state anxiety. *Complementary Therapies in Medicine*, 21(5), pp.468-72.

Novosad C and Thoman E B. (2003). The breathing bear: an intervention for crying babies and their mothers. *Journal of Developmental & Behavioral Pediatrics*, 24(2), pp.89-95.

Nyatsanza M, Schneider M and Davies T ; Lund C ;. (2016). Filling the treatment gap: developing a task sharing counselling intervention for perinatal depression in Khayelitsha, South Africa. *BMC Psychiatry*, 16, pp.164.

O'Hara M W, Pearlstein T and Stuart S ; Long J D; Mills J A; Zlotnick C ;. (2019). A placebo controlled treatment trial of sertraline and interpersonal psychotherapy for postpartum depression. *Journal of Affective Disorders*, 245, pp.524-532.

O'Mahen H A and Flynn H A;. (2008). Preferences and perceived barriers to treatment for depression during the perinatal period. *Journal of Women's Health*, 17(8), pp.1301-9.

O'Mahen H A, Woodford J and McGinley J ; Warren F C; Richards D A; Lynch T R; Taylor R S;. (2013). Internet-based behavioral activation--treatment for postnatal depression (Netmums): a randomized controlled trial. *Journal of Affective Disorders*, 150(3), pp.814-22.

O'Mahen H, Himle J A and Fedock G ; Henshaw E ; Flynn H ;. (2013). A pilot randomized controlled trial of cognitive behavioral therapy for perinatal depression adapted for women with low incomes. *Depression & Anxiety*, 30(7), pp.679-87.

O'Mahen H A, Richards D A; Woodford J and Wilkinson E ; McGinley J ; Taylor R S; Warren F C;. (2014). Netmums: a phase II randomized controlled trial of a guided Internet behavioural activation treatment for postpartum depression. *Psychological Medicine*, 44(8), pp.1675-89.

Okronipa H, Adu-Afarwuah S and Lartey A ; Ashorn P ; Vosti S A; Young R R; Dewey K G;. (2018). Maternal supplementation with small-quantity lipid-based nutrient supplements during pregnancy and lactation does not reduce depressive symptoms at 6 months postpartum in Ghanaian women: a randomized controlled trial. *Archives of Women's Mental Health*, 21(1), pp.55-63.

Olayiwola J N, Irizarry O C; O'Connell K and Milan S ;. (2013). Living Smart, Living Fit: a patient-centered program to improve perinatal outcomes in a community health center population. *Journal of Primary Care & Community Health*, 4(1), pp.31-5.

Olhaberry M, Leon M J and Seguel M ; Mena C ;. (2015). Video-feedback intervention in mother-baby dyads with depressive symptomatology and relationship difficulties. *Research in Psychotherapy: Psychopathology and Process and Outcome*, 18(2), pp.82-92.

Onozawa K, Glover V and Adams D ; Modi N ; Kumar R C;. (2001). Infant massage improves mother-infant interaction for mothers with postnatal depression. *Journal of Affective Disorders*, 63(1-3), pp.201-7.

Orengo-Aguayo R E and Segre L S;. (2016). Depression treatment delivered at the point-of-care: a qualitative assessment of the views of low-income US mothers. *Journal of Reproductive & Infant Psychology*, 34(1), pp.35-48.

Ormsby S M and Dahlen H G; Smith C A;. (2018). Women's experiences of having depression during pregnancy and receiving acupuncture treatment-A qualitative study. *Women and Birth*, 31(6), pp.469-478.

- Ortiz Collado, M A and Saez M ; Favrod J ; Hatem M ;. (2014). Antenatal psychosomatic programming to reduce postpartum depression risk and improve childbirth outcomes: a randomized controlled trial in Spain and France. *BMC Pregnancy & Childbirth*, 14, pp.22.
- Oswalt K L and McClain D B; Melnyk B. (2013). Reducing anxiety among children born preterm and their young mothers. *MCN and American Journal of Maternal Child Nursing*, 38(3), pp.144-9.
- Pan W L and Chang C W; Chen S M; Gau M L;. (2019). Assessing the effectiveness of mindfulness-based programs on mental health during pregnancy and early motherhood - a randomized control trial. *BMC Pregnancy & Childbirth*, 19(1), pp.346.
- Panyayong B. (2013). Postpartum depression among Thai women: a national survey. *Journal of the Medical Association of Thailand*, 96(7), pp.761-7.
- Paris R, Bolton R E and Spielman E ;. (2011). Evaluating a home-based dyadic intervention: Changes in postpartum depression, maternal perceptions, and mother-infant interactions. *Infant Mental Health Journal*, 32(3), pp.319-338.
- Parker G B, Hegarty B and Paterson A ; Hadzi-Pavlovic D ; Granville-Smith I ; Gokiert A ;. (2015). Predictors of post-natal depression are shaped distinctly by the measure of 'depression'. *Journal of Affective Disorders*, 173, pp.239-44.
- Peindl K S and Wisner K L; Hanusa B H;. (2004). Identifying depression in the first postpartum year: guidelines for office-based screening and referral. *Journal of Affective Disorders*, 80(1), pp.37-44.
- Perez-Blasco J, Viguer P and Rodrigo M F;. (2013). Effects of a mindfulness-based intervention on psychological distress, well-being, and maternal self-efficacy in breast-feeding mothers: results of a pilot study. *Archives of Women's Mental Health*, 16(3), pp.227-36.
- Petch S, DeMaio A and Daly S ;. (2019). Prediction of recurrent preterm delivery in asymptomatic women- an anxiety reducing measure?. *European Journal of Obstetrics & Gynecology and Reproductive Biology*, 4, pp.100064.
- Phua D Y, Kee Mzkl and Koh D X. P; Rifkin-Graboi A ; Daniels M ; Chen H ; Chong Y S; Broekman B F. P; Magiati I ; Karnani N ; Pluess M ; Meaney M J; Growing Up In Singapore Towards Healthy Outcomes Study; Group ;. (2017). Positive maternal mental health during pregnancy associated with specific forms of adaptive development in early childhood: Evidence from a longitudinal study. *Development & Psychopathology*, 29(5), pp.1573-1587.
- Polman R, Kaiseler M and Borkoles E ;. (2007). Effect of a single bout of exercise on the mood of pregnant women. *Journal of Sports Medicine and Physical Fitness*, 47(1), pp.103-111.
- Posmontier B, Neugebauer R and Stuart S ; Chittams J ; Shaughnessy R ;. (2016). Telephone-Administered Interpersonal Psychotherapy by Nurse-Midwives for Postpartum Depression. *Journal of Midwifery & Women's Health*, 61(4), pp.456-66.
- Posmontier Bobbie, Bina Rena and Glasser Saralee ; Cinamon Tali ; Styr Benedict ; Sammarco Therese ;. (2019). Incorporating Interpersonal Psychotherapy for Postpartum

Depression Into Social Work Practice in Israel. *Research on Social Work Practice*, 29(1), pp.61-68.

Powell C, Bamber D and Long J ; Garratt R ; Brown J ; Rudge S ; Morris T ; Bhupendra Jaicim ; N ; Plachcinski R ; Dyson S ; Boyle E M; St James-Roberts ; I ;. (2018). Mental health and well-being in parents of excessively crying infants: Prospective evaluation of a support package. *Child: Care and Health & Development*, 44(4), pp.607-615.

Premji S, McDonald S W and Metcalfe A ; Faris P ; Quan H ; Tough S ; McNeil D A;. (2019). Examining postpartum depression screening effectiveness in well child clinics in Alberta, Canada: A study using the All Our Families cohort and administrative data. *Preventive Medicine Reports*, 14, pp.100888.

Prevatt B S and Lowder E M; Desmarais S L;. (2018). Peer-support intervention for postpartum depression: Participant satisfaction and program effectiveness. *Midwifery*, 64, pp.38-47.

Prevatt B S and Desmarais S L;. (2018). Facilitators and Barriers to Disclosure of Postpartum Mood Disorder Symptoms to a Healthcare Provider. *Maternal & Child Health Journal*, 22(1), pp.120-129.

Preyde M and Ardal F . (2003). Effectiveness of a parent "buddy" program for mothers of very preterm infants in a neonatal intensive care unit. *CMAJ Canadian Medical Association Journal*, 168(8), pp.969-73.

Price C, Kantrowitz-Gordon I and Calhoun R ;. (2019). A pilot feasibility study of mindfulness childbirth education for women with a history of sexual trauma. *Complementary Therapies in Clinical Practice*, 37, pp.102-108.

Priest S R, Henderson J and Evans S F; Hagan R ;. (2003). Stress debriefing after childbirth: a randomised controlled trial. *Medical Journal of Australia*, 178(11), pp.542-5.

Pritchett R, Jolly K and Daley A J; Turner K ; Bradbury-Jones C ; Team P AM-PeRS Study;. (2017). Women's experiences of exercise as a treatment for their postnatal depression: A nested qualitative study. *Journal of Health Psychology*, , pp.1359105317726590.

Pugh N E and Hadjistavropoulos H D; Dirkse D. (2016). A Randomised Controlled Trial of Therapist-Assisted, Internet-Delivered Cognitive Behavior Therapy for Women with Maternal Depression. *PLoS ONE [Electronic Resource]*, 11(3), pp.e0149186.

Puryear L J, Nong Y H; Correa N P; Cox K and Greeley C S;. (2019). Outcomes of Implementing Routine Screening and Referrals for Perinatal Mood Disorders in an Integrated Multi-site Pediatric and Obstetric Setting. *Maternal & Child Health Journal*, 23(10), pp.1292-1298.

Quispel C, Lambregtse-van den Berg and M ; Kaan M ; van den Berg ; P M ; Hoogendijk W J; Steegers E A; Koudstaal T ; Muis I ; Bonsel G J;. (2014). Withdrawal from mental and psychosocial care during pregnancy. *Journal of Psychosomatic Obstetrics & Gynecology*, 35(4), pp.140-5.

Rabiei L, Mazaheri M A and Masoudi R ; Hasheminia S A;. (2014). Fordyce happiness program and postpartum depression. *Journal of Research in Medical Sciences*, 19(3), pp.251-6.



Rahman A. (2007). Challenges and opportunities in developing a psychological intervention for perinatal depression in rural Pakistan--a multi-method study. *Archives of Women's Mental Health*, 10(5), pp.211-9.

Ramos-Marcuse F, Oberlander S E and Papas M A; McNary S W; Hurley K M; Black M M;. (2010). Stability of maternal depressive symptoms among urban, low-income, African American adolescent mothers. *Journal of Affective Disorders*, 122(1-2), pp.68-75.

Ramphos E S and Kelman A R; Stanley M L; Barrera A Z;. (2019). Responding to women's needs and preferences in an online program to prevent postpartum depression. *Internet Interventions*, 18 (no pagination), pp..

Reay R, Stuart S and Owen C ;. (2003). Implementation and effectiveness of interpersonal psychotherapy in a community mental health service. *Australasian Psychiatry*, 11(3), pp.284-289.

Reay R, Fisher Y and Robertson M ; Adams E ; Owen C ; Kumar R ;. (2006). Group interpersonal psychotherapy for postnatal depression: a pilot study. *Archives of Women's Mental Health*, 9(1), pp.31-9.

Reid M, Glazener C and Murray G D; Taylor G S;. (2002). A two-centred pragmatic randomised controlled trial of two interventions of postnatal support. *BJOG: An International Journal of Obstetrics & Gynaecology*, 109(10), pp.1164-70.

Reilly N, Brake E and Kalra H ; Austin M P;. (2019). Insights into implementation of routine depression screening and psychosocial assessment in a private hospital setting: A qualitative study. *Australian & New Zealand Journal of Obstetrics & Gynaecology*, 24, pp.24.

Reinstein S, Lieb K and Bernstein P S; Karkowsky C E;. (2019). The best perinatal depression screening: Is self-administered PHQ2 more feasible than a nurse-administered one?. *Perspectives in Psychiatric Care*, 18, pp.18.

Reupert A and Maybery D . (2011). Programmes for parents with a mental illness. *Journal of Psychiatric & Mental Health Nursing*, 18(3), pp.257-64.

Ribeiro M K. A, Alcantara-Silva T R. M; Oliveira J C. M; Paula T C; Dutra J B. R; Pedrino G R; Simoes K and Sousa R B; Rebelo A C. S;. (2018). Music therapy intervention in cardiac autonomic modulation, anxiety, and depression in mothers of preterms: randomized controlled trial. *BMC psychology*, 6(1), pp.57.

Robichaud Karen Petty. (2009). The effects of an exercise intervention on the psychological well-being of postpartum women. *Dissertation Abstracts International: Section B: The Sciences and Engineering*, 69(7-B), pp.4124.

Robinson S and Young J . (1982). Screening for depression and anxiety in the post-natal period: acceptance or rejection of a subsequent treatment offer. *Australian & New Zealand Journal of Psychiatry*, 16(2), pp.47-51.

Rock M M. (2019). A Stepped-Collaborative Perinatal Depression Model. *Journal of the American Psychiatric Nurses Association*, 25(5), pp.376-384.

Rollans M, Schmied V and Kemp L ; Meade T ;. (2013). 'We just ask some questions...' the process of antenatal psychosocial assessment by midwives. *Midwifery*, 29(8), pp.935-42.

- Rollans M, Kohlhoff J and Meade T ; Kemp L ; Schmieid V ;. (2016). Partner Involvement: Negotiating the Presence of Partners in Psychosocial Assessment as Conducted by Midwives and Child and Family Health Nurses. *Infant Mental Health Journal*, 37(3), pp.302-12.
- Rossiter C, Fowler C and McMahon C ; Kowalenko N ;. (2012). Supporting depressed mothers at home: their views on an innovative relationship-based intervention. *Contemporary Nurse*, 41(1), pp.90-100.
- Rotheram-Fuller E J, Tomlinson M and Scheffler A ; Weichle T W; Hayati Rezvan ; P ; Comulada W S; Rotheram-Borus M J;. (2018). Maternal patterns of antenatal and postnatal depressed mood and the impact on child health at 3-years postpartum. *Journal of Consulting & Clinical Psychology*, 86(3), pp.218-230.
- Rowan P, Greisinger A and Brehm B ; Smith F ; McReynolds E ;. (2012). Outcomes from implementing systematic antepartum depression screening in obstetrics. *Archives of Women's Mental Health*, 15(2), pp.115-20.
- Rowe H J and Fisher J R;. (2010). The contribution of Australian residential early parenting centres to comprehensive mental health care for mothers of infants: evidence from a prospective study. *International Journal of Mental Health Systems*, 4, pp.6.
- Rowe H J and Wynter K H; Burns J K; Fisher J R. W;. (2017). A complex postnatal mental health intervention: Australian translational formative evaluation. *Health Promotion International*, 32(4), pp.610-623.
- Ruiz R J, Newman M and Records K ; Wommack J C; Stowe R P; Pasillas R M;. (2019). Pilot Study of the Mastery Lifestyle Intervention. *Nursing Research*, 06, pp.06.
- Russomagno S and Waldrop J . (2019). Improving Postpartum Depression Screening and Referral in Pediatric Primary Care. *Journal of Pediatric Health Care*, 33(4), pp.e19-e27.
- Saenz P, Cerda M and Diaz J L; Yi P ; Gorba M ; Boronat N ; Barreto P ; Vento M ;. (2009). Psychological stress of parents of preterm infants enrolled in an early discharge programme from the neonatal intensive care unit: a prospective randomised trial. *Archives of Disease in Childhood Fetal & Neonatal Edition*, 94(2), pp.F98-F104.
- Salonen A H and Pridham K F; Brown R L; Kaunonen M. (2014). Impact of an internet-based intervention on Finnish mothers' perceptions of parenting satisfaction, infant centrality and depressive symptoms during the postpartum year. *Midwifery*, 30(1), pp.112-22.
- Sampson McClain, Villarreal Yolanda and Rubin Allen ;. (2016). A problem-solving therapy intervention for low-income, pregnant women at risk for postpartum depression. *Research on Social Work Practice*, 26(3), pp.236-242.
- San Martin Porter, M A and Betts K ; Kisely S ; Pecoraro G ; Alati R ;. (2019). Screening for perinatal depression and predictors of underscreening: findings of the Born in Queensland study. *Medical Journal of Australia*, 210(1), pp.32-37.
- Sawyer A, Kaim A and Le H N; McDonald D ; Mittinty M ; Lynch J ; Sawyer M ;. (2019). The Effectiveness of an App-Based Nurse-Moderated Program for New Mothers With Depression and Parenting Problems (eMums Plus): Pragmatic Randomized Controlled Trial. *Journal of medical Internet research*, 21(6), pp.e13689.

Schaar G L and Hall M. (2013). A nurse-led initiative to improve obstetricians' screening for postpartum depression. *Nursing for Women's Health*, 17(4), pp.306-15; quiz 316.

Schmied V, Langdon R and Matthey S ; Kemp L ; Austin M P; Johnson M ;. (2016). Antenatal psychosocial risk status and Australian women's use of primary care and specialist mental health services in the year after birth: a prospective study. *BMC Women's Health*, 16(1), pp.69.

Scholle S H and Haskett R F; Hanusa B H; Pincus H A; Kupfer D J;. (2003). Addressing depression in obstetrics/gynecology practice. *General Hospital Psychiatry*, 25(2), pp.83-90.

Seeley J R, Sheeber L B; Feil E G; Leve C and Davis B ; Sorensen E ; Allan S ;. (2019). Mediation analyses of Internet-facilitated cognitive behavioral intervention for maternal depression. *Cognitive behaviour therapy*, 48(4), pp.337-352.

Segre L S and Stasik S M; O'Hara M W; Arndt S. (2010). Listening visits: an evaluation of the effectiveness and acceptability of a home-based depression treatment. *Psychotherapy Research*, 20(6), pp.712-21.

Segre L S and O'Hara M W; Brock R L; Taylor D. (2012). Depression screening of perinatal women by the Des Moines Healthy Start Project: program description and evaluation. *Psychiatric Services*, 63(3), pp.250-5.

Segre L S and O'Hara M W; Fisher S D;. (2013). Perinatal depression screening in healthy start: an evaluation of the acceptability of technical assistance consultation. *Community Mental Health Journal*, 49(4), pp.407-11.

Segre L S and Brock R L; O'Hara M W;. (2015). Depression treatment for impoverished mothers by point-of-care providers: A randomized controlled trial. *Journal of Consulting & Clinical Psychology*, 83(2), pp.314-24.

Segre L S and Orengo-Aguayo R E; Siewert R C;. (2016). Depression Management by NICU Nurses: Mothers' Views. *Clinical Nursing Research*, 25(3), pp.273-90.

Selkirk R, McLaren S and Ollerenshaw A ; McLachlan A J; Moten J ;. (2006). The longitudinal effects of midwife-led postnatal debriefing on the psychological health of mothers. *Journal of Reproductive & Infant Psychology*, 24(2), pp.133-147.

Sezen C and Unsalver B O. (2019). Group art therapy for the management of fear of childbirth. *Arts in Psychotherapy*, 64, pp.9-19.

Shahnazi H, Sabooteh S and Sharifirad G ; Mirkarimi K ; Hassanzadeh A ;. (2015). The impact of education intervention on the Health Belief Model constructs regarding anxiety of nulliparous pregnant women. *Journal of Education & Health Promotion*, 4, pp.27.

Shahtaheri E, Hosseinian S and Ebrahimi M ; Jalilabadi Z ; Mirderkvand F ;. (2016). The impact of stress reduction program based on mindfulness and conscious yoga on depression, perceived stress and quality of life in pregnant women. *Acta Medica Mediterranea*, 32(SpecialIssue5), pp.2047-2050.

Sharan H, Kaplan B and Weizer N ; Sulkes J ; Merlob P ;. (2006). Early screening of postpartum depression using the Edinburgh Postnatal Depression Scale. *International Journal of Risk and Safety in Medicine*, 18(4), pp.213-218.

- Sharma V, Khan M and Corpse C ; Sharma P ;. (2008). Missed bipolarity and psychiatric comorbidity in women with postpartum depression. *Bipolar Disorders*, 10(6), pp.742-7.
- Sharp D J, Chew-Graham C and Tylee A ; Lewis G ; Howard L ; Anderson I ; Abel K ; Turner K M; Hollinghurst S P; Tallon D ; McCarthy A ; Peters T J;. (2010). A pragmatic randomised controlled trial to compare antidepressants with a community-based psychosocial intervention for the treatment of women with postnatal depression: the RESPOND trial. *Health Technology Assessment (Winchester and England)*, 14(43), pp.iii-iv, ix-xi, 1-153.
- Shaw R J, St John and N ; Lilo E A; Jo B ; Benitz W ; Stevenson D K; Horwitz S M;. (2013). Prevention of traumatic stress in mothers with preterm infants: a randomized controlled trial. *Pediatrics*, 132(4), pp.e886-94.
- Sheeber L B, Feil E G; Seeley J R; Leve C and Gau J M; Davis B ; Sorensen E ; Allan S ;. (2017). Mom-net: Evaluation of an internet-facilitated cognitive behavioral intervention for low-income depressed mothers. *Journal of Consulting and Clinical Psychology*, 85(4), pp.355-366.
- Sheeder J, Kabir K and Stafford B ;. (2009). Screening for postpartum depression at well-child visits: is once enough during the first 6 months of life?. *Pediatrics*, 123(6), pp.e982-8.
- Shenai N, Gopalan P and Glance J ;. (2019). Integrated Brief Intervention for PTSD and Substance Use in an Antepartum Unit. *Maternal & Child Health Journal*, 23(5), pp.592-596.
- Sherman L J and Ali M M;. (2018). Diagnosis of Postpartum Depression and Timing and Types of Treatment Received Differ for Women with Private and Medicaid Coverage. *Women's Health Issues*, 28(6), pp.524-529.
- Shields N, Reid M and Cheyne H ; Holmes A ; McGinley M ; Turnbull D ; Smith L N;. (1997). Impact of midwife-managed care in the postnatal period: An exploration of psychosocial outcomes. *Journal of Reproductive and Infant Psychology*, 15(2), pp.91-108.
- Shivakumar G, Brandon A R and Johnson N L; Freeman M P;. (2014). Screening to Treatment: Obstacles and Predictors in Perinatal Depression (STOP-PPD) in the Dallas Healthy Start program. *Archives of Women's Mental Health*, 17(6), pp.575-8.
- Shorey Shefaly. (2013). The effectiveness of a Postnatal Psychoeducation Programme on outcomes of first-time mothers in Singapore. *Effectiveness of a Postnatal Psychoeducation Programme on Outcomes of First-time Mothers in Singapore*, , pp.1-1.
- Shorey S, Chan S W and Chong Y S; He H G;. (2015). A randomized controlled trial of the effectiveness of a postnatal psychoeducation programme on self-efficacy, social support and postnatal depression among primiparas. *Journal of Advanced Nursing*, 71(6), pp.1260-73.
- Shulman B, Dueck R and Ryan D ; Breau G ; Sadowski I ; Misri S ;. (2018). Feasibility of a mindfulness-based cognitive therapy group intervention as an adjunctive treatment for postpartum depression and anxiety. *Journal of Affective Disorders*, 235, pp.61-67.
- Sikander Siham, Ahmad Ikhlaiq and Atif Najia ; Zaidi Ahmed ; Vanobberghen Fiona ; Weiss Helen A; Nisar Anum ; Tabana Hanani ; Ain Qurat U; Bibi Amina ; Bilal Samina ; Bibi Tayyiba ; Liaqat Rakshanda ; Sharif Maria ; Zulfiqar Shaffaq ; Fuhr Daniela C; Price LeShawndra N; Patel Vikram ; Rahman Atif ;. (2019). Delivering the Thinking Healthy

Programme for perinatal depression through volunteer peers: A cluster randomised controlled trial in Pakistan. *The Lancet Psychiatry*, 6(2), pp.128-139.

Silverstein M, Feinberg E and Cabral H ; Sauder S ; Egbert L ; Schainker E ; Kamholz K ; Hegel M ; Beardslee W ;. (2011). Problem-solving education to prevent depression among low-income mothers of preterm infants: a randomized controlled pilot trial. *Archives of Women's Mental Health*, 14(4), pp.317-24.

Silverstein M, Diaz-Linhart Y and Cabral H ; Beardslee W ; Broder-Fingert S ; Kistin C J ; Patts G ; Feinberg E ;. (2018). Engaging mothers with depressive symptoms in care: Results of a randomized controlled trial in head start. *Psychiatric Services*, 69(11), pp.1175-1180.

Simavli S, Kaygusuz I and Gumus I ; Usluogullari B ; Yildirim M ; Kafali H ;. (2014). Effect of music therapy during vaginal delivery on postpartum pain relief and mental health. *Journal of Affective Disorders*, 156, pp.194-9.

Simeone S, Lojo C and Garcia-Esteve L ; Triunfo S ; Crovetto F ; Arranz A ; Gratacos E ; Figueras F ;. (2015). Psychological impact of first-trimester prevention for preeclampsia on anxiety. *Prenatal Diagnosis*, 35(1), pp.60-4.

Siu B W. M and Chow H M; Kwok S S; Li O L; Koo M L; Poon P W;. (2011). Pathway to care for postnatal depression after implementation of the Comprehensive Child Development Service in a regional hospital in Hong Kong. *Asia-Pacific Psychiatry*, 3(2), pp.54-60.

Skoog M, Berggren V and Hallstrom I K;. (2019). 'Happy that someone cared'-Non-native-speaking immigrant mothers' experiences of participating in screening for postpartum depression in the Swedish child health services. *Journal of Child Health Care*, 23(1), pp.118-130.

Small Rhonda, Lumley Judith and Donohue Lisa ; Potter Anne ; Waldenstrom Ulla ;. (2000). Randomised controlled trial of midwife led debriefing to reduce maternal depression after operative childbirth. *BMJ: British Medical Journal*, 321(7268), pp.1043-1047.

Smith M V, Rosenheck R A; Cavaleri M A; Howell H B; Poschman K and Yonkers K A;. (2004). Screening for and detection of depression, panic disorder, and PTSD in public-sector obstetric clinics. *Psychiatric Services*, 55(4), pp.407-14.

Smith Sandra and Moore Elizabeth . (2012). Health Literacy and Depression in the Context of Home Visitation. *Maternal & Child Health Journal*, 16(7), pp.1500-1508.

Smith Sarah Courtney. (2018). Evaluating a stepped care protocol for postpartum depression in a pediatric primary care clinic. *Dissertation Abstracts International: Section B: The Sciences and Engineering*, 79(4-B(E)), pp.No Pagination Specified.

SmithBattle L, Chantamit O Pas C and Freed P ; McLaughlin D ; Schneider J K;. (2017). Moms growing together: Piloting action methods and expressive arts in a therapeutic group for teen mothers. *Journal of Child & Adolescent Psychiatric Nursing*, 30(2), pp.72-79.

Sorg M, Coddington J and Ahmed A ; Richards E ;. (2019). Improving Postpartum Depression Screening in Pediatric Primary Care: A Quality Improvement Project. *Journal of Pediatric Nursing*, 46, pp.83-88.

Sorsdahl K, Petersen Williams and P ; Everett-Murphy K ; Vythilingum B ; de Villiers ; P ; Myers B ; Stein D J;. (2015). Feasibility and Preliminary Responses to a Screening and Brief Intervention Program for Maternal Mental Disorders Within the Context of Primary Care. *Community Mental Health Journal*, 51(8), pp.962-9.

Steinberg S I and Bellavance F. (1999). Characteristics and treatment of women with antenatal and postpartum depression. *International Journal of Psychiatry in Medicine*, 29(2), pp.209-33.

Stevens N R, Heath N M; Lillis T A; McMinn K and Tirone V ; Sha'ini M ;. (2018). Examining the effectiveness of a coordinated perinatal mental health care model using an intersectional-feminist perspective. *Journal of behavioral medicine*, 41(5), pp.627-640.

Stewart R C, Ashorn P and Umar E ; Dewey K G; Ashorn U ; Creed F ; Rahman A ; Tomenson B ; Prado E L; Maleta K ;. (2017). The impact of maternal diet fortification with lipid-based nutrient supplements on postpartum depression in rural Malawi: a randomised-controlled trial. *Maternal & Child Nutrition*, 13(2), pp.04.

Stramrood C A, van der Velde and J ; Doornbos B ; Marieke Paarlberg ; K ; Weijmar Schultz ; W C ; van Pampus ; M G ;. (2012). The patient observer: eye-movement desensitization and reprocessing for the treatment of posttraumatic stress following childbirth. *Birth (Berkeley and Calif.)*, 39(1), pp.70-76.

Surjaningrum E R, Jorm A F; Minas H and Kakuma R ;. (2018). Personal attributes and competencies required by community health workers for a role in integrated mental health care for perinatal depression: voices of primary health care stakeholders from Surabaya, Indonesia. *International Journal of Mental Health Systems*, 12, pp.46.

Surkan P J, Gottlieb B R; McCormick M C; Hunt A and Peterson K E;. (2012). Impact of a health promotion intervention on maternal depressive symptoms at 15 months postpartum. *Maternal & Child Health Journal*, 16(1), pp.139-48.

Swanson L M, Burgess H J; Zollars J and Todd Arnedt ; J ;. (2018). An open-label pilot study of a home wearable light therapy device for postpartum depression. *Archives of Women's Mental Health*, 21(5), pp.583-586.

Symon B and Crichton G E. (2017). The joy of parenting: infant sleep intervention to improve maternal emotional well-being and infant sleep. *Singapore Medical Journal*, 58(1), pp.50-54.

Ta Park, V M and Goyal D ; Nguyen T ; Lien H ; Rosidi D ;. (2017). Postpartum Traditions, Mental Health, and Help-Seeking Considerations Among Vietnamese American Women: a Mixed-Methods Pilot Study. *Journal of Behavioral Health Services & Research*, 44(3), pp.428-441.

Tabeshpour J, Sobhani F and Sadjadi S A; Hosseinzadeh H ; Mohajeri S A; Rajabi O ; Taherzadeh Z ; Eslami S ;. (2017). A double-blind, randomized, placebo-controlled trial of saffron stigma (*Crocus sativus* L.) in mothers suffering from mild-to-moderate postpartum depression. *Phytomedicine*, 36, pp.145-152.

Tabor A and Jonsson M H. (1987). Psychological impact of amniocentesis on low-risk women. *Prenatal Diagnosis*, 7(6), pp.443-449.

Tachibana Y, Koizumi N and Akanuma C ; Tarui H ; Ishii E ; Hoshina T ; Suzuki A ; Asano A ; Sekino S ; Ito H ;. (2019). Integrated mental health care in a multidisciplinary maternal and child health service in the community: the findings from the Suzaka trial. *BMC Pregnancy & Childbirth*, 19(1), pp.58.

Tandon S D, Parillo K M; Jenkins C and Duggan A K;. (2005). Formative evaluation of home visitors' role in addressing poor mental health, domestic violence, and substance abuse among low-income pregnant and parenting women. *Maternal & Child Health Journal*, 9(3), pp.273-283.

Tandon S D, Perry D F; Mendelson T and Kemp K ; Leis J A;. (2011). Preventing perinatal depression in low-income home visiting clients: a randomized controlled trial. *Journal of Consulting & Clinical Psychology*, 79(5), pp.707-12.

Teissedre F and Chabol H . (2003). Postnatal depression: a study of the predictive effects of postnatal anxiety. *Irish Journal of Psychological Medicine*, 20(4), pp.111-114.

Teychenne M, van der Pligt and P ; Abbott G ; Brennan L ; Olander E K;. (2018). Feasibility and acceptability of a home-based physical activity program for postnatal women with depressive symptoms: A pilot study. *Mental Health and Physical Activity*, 14, pp.82-89.

Thippeswamy H, Desai G and Chandra P ;. (2018). Help-seeking patterns in women with postpartum severe mental illness: a report from southern India. *Archives of Women's Mental Health*, 21(5), pp.573-578.

Thitipitchayanant K, Somrongthong R and Kumar R ; Kanchanakharn N ;. (2018). Effectiveness of self-empowerment-affirmation-relaxation(Self-EAR) program for postpartum blues mothers: A randomized controlled trial. *Pakistan Journal of Medical Sciences*, 34(6), pp.1488-1493.

Thoppil J, Riutcel T L and Nalesnik S W;. (2005). Early intervention for perinatal depression. *American Journal of Obstetrics & Gynecology*, 192(5), pp.1446-8.

Thorpe Karen. (1993). A study of the use of the Edinburgh Postnatal Depression Scale with parent groups outside the postpartum period. *Journal of Reproductive and Infant Psychology*, 11(2), pp.119-125.

Timpano K R and Abramowitz J S; Mahaffey B L; Mitchell M A; Schmidt N B;. (2011). Efficacy of a prevention program for postpartum obsessive-compulsive symptoms. *Journal of Psychiatric Research*, 45(11), pp.1511-7.

Toler S, Stapleton S and Kertsburg K ; Callahan T J; Hastings-Tolsma M ;. (2018). Screening for postpartum anxiety: A quality improvement project to promote the screening of women suffering in silence. *Midwifery*, 62, pp.161-170.

Tragea C, Chrousos G P and Alexopoulos E C; Darviri C ;. (2014). A randomized controlled trial of the effects of a stress management programme during pregnancy. *Complementary Therapies in Medicine*, 22(2), pp.203-11.

Traube Renee. (2017). Development of a Quality Improvement Initiative to Screen for Postpartum Depression. *Development of a Quality Improvement Initiative to Screen for Postpartum Depression*, , pp.1-1.

Tripathy P, Nair N and Barnett S ; Mahapatra R ; Borghi J ; Rath S ; Rath S ; Gope R ; Mahto D ; Sinha R ; Lakshminarayana R ; Patel V ; Pagel C ; Prost A ; Costello A ;. (2010). Effect of a participatory intervention with women's groups on birth outcomes and maternal depression in Jharkhand and Orissa, India: a cluster-randomised controlled trial. *Lancet*, 375(9721), pp.1182-92.

Tsai A C, Tomlinson M and Dewing S ; le Roux ; I M ; Harwood J M; Chopra M ; Rotheram-Borus M J;. (2014). Antenatal depression case finding by community health workers in South Africa: feasibility of a mobile phone application. *Archives of Women's Mental Health*, 17(5), pp.423-31.

Tubay A T, Mansalis K A; Simpson M J; Armitage N H; Briscoe G and Potts V ;. (2019). The Effects of Group Prenatal Care on Infant Birthweight and Maternal Well-Being: A Randomized Controlled Trial. *Military Medicine*, 184(5-6), pp.e440-e446.

Turkstra E, Mihala G and Scuffham P A; Creedy D K; Gamble J ; Toohill J ; Fenwick J ;. (2017). An economic evaluation alongside a randomised controlled trial on psycho-education counselling intervention offered by midwives to address women's fear of childbirth in Australia. *Sexual & reproductive healthcare : official journal of the Swedish Association of Midwives*, 11, pp.1-6.

Uebelacker Lisa, Battle Cynthia and Sutton Kaeli ; Magee Susanna ; Miller Ivan ;. (2016). A pilot randomized controlled trial comparing prenatal yoga to perinatal health education for antenatal depression. *Archives of Women's Mental Health*, 19(3), pp.543-547.

Ugarriza D N. (2004). Group therapy and its barriers for women suffering from postpartum depression. *Archives of Psychiatric Nursing*, 18(2), pp.39-48.

Ugarriza D N and Schmidt L. (2006). Telecare for women with postpartum depression. *Journal of Psychosocial Nursing & Mental Health Services*, 44(1), pp.37-45.

Upshur C C, Wenz-Gross M and Weinreb L ; Moffitt J J;. (2016). Using Prenatal Advocates to Implement a Psychosocial Education Intervention for Posttraumatic Stress Disorder during Pregnancy: Feasibility, Care Engagement, and Predelivery Behavioral Outcomes. *Womens Health Issues*, 26(5), pp.537-45.

van der Ham , J and Berry K ; Hoehn E ; Fraser J ;. (2013). A collaborative approach to perinatal and infant mental health service delivery in Australia. *Australasian Psychiatry*, 21(4), pp.371-5.

van der Zee-van den Berg, A I and Boere-Boonekamp M M; Groothuis-Oudshoorn C G. M; MJ I Jzerman; Haasnoot-Smallegange R M. E; Reijneveld S A;. (2017). Post-Up Study: Postpartum Depression Screening in Well-Child Care and Maternal Outcomes. *Pediatrics*, 140(4), pp..

Van Lieshout , R J and Layton H ; Feller A ; Ferro M A; Biscaro A ; Bieling P J;. (2019). Public health nurse delivered group cognitive behavioral therapy (CBT) for postpartum depression: A pilot study. *Public Health Nursing*, 15, pp.15.

Van Ravesteyn , L M and Kamperman A M; Schneider T A. J; Raats M E; Steegers E A. P; Tiemeier H ; Hoogendijk W J. G; Lambregtse-van den Berg; M P ;. (2018). Group-based multicomponent treatment to reduce depressive symptoms in women with co-morbid



- psychiatric and psychosocial problems during pregnancy: A randomized controlled trial. *Journal of Affective Disorders*, 226, pp.36-44.
- Vargas-Terrones M, Barakat R and Santacruz B ; Fernandez-Buhigas I ; Mottola M F;. (2019). Physical exercise programme during pregnancy decreases perinatal depression risk: a randomised controlled trial. *British Journal of Sports Medicine*, 53(6), pp.348-353.
- Venkatesh K K, Nadel H and Blewett D ; Freeman M P; Kaimal A J; Riley L E;. (2016). Implementation of universal screening for depression during pregnancy: feasibility and impact on obstetric care. *American Journal of Obstetrics & Gynecology*, 215(4), pp.517.e1-8.
- Verdeli H, Therosme T and Eustache E ; Hilaire O S; Joseph B ; Sonmez C C; Raviola G ;. (2016). Community Norms and Human Rights: Supervising Haitian Colleagues on Interpersonal Psychotherapy (IPT) With a Depressed and Abused Pregnant Woman. *Journal of Clinical Psychology*, 72(8), pp.847-55.
- Victoria C G, Langer A and Barros F ; Belizan J ; Farnot U ; Villar J ;. (1994). The Latin American Multicenter Trial on psychosocial support during pregnancy: methodology and baseline comparability. Latin American Network for Perinatal and Reproductive Research (LANPER). *Controlled Clinical Trials*, 15(5), pp.379-94.
- Vieten C and Astin J . (2008). Effects of a mindfulness-based intervention during pregnancy on prenatal stress and mood: results of a pilot study. *Archives of Women's Mental Health*, 11(1), pp.67-74.
- Vieten C, Laraia B A and Kristeller J ; Adler N ; Coleman-Phox K ; Bush N R; Wahbeh H ; Duncan L G; Epel E ;. (2018). The mindful moms training: development of a mindfulness-based intervention to reduce stress and overeating during pregnancy. *BMC Pregnancy & Childbirth*, 18(1), pp.201.
- Vigod S N and Murphy K E; Dennis C L; Oberlander T F; Ray J G; Daskalakis Z J; Blumberger D M;. (2019). Transcranial direct current stimulation (tDCS) for depression in pregnancy: A pilot randomized controlled trial. *Brain Stimulation*, 12(6), pp.1475-1483.
- Vigod S N, Hussain-Shamsy N and Stewart D E; Grigoriadis S ; Metcalfe K ; Oberlander T F; Schram C ; Taylor V H; Dennis C L;. (2019). A patient decision aid for antidepressant use in pregnancy: Pilot randomized controlled trial. *Journal of Affective Disorders*, 251, pp.91-99.
- Viveiros C J and Darling E K;. (2018). Barriers and facilitators of accessing perinatal mental health services: The perspectives of women receiving continuity of care midwifery. *Midwifery*, 65, pp.8-15.
- Wahlbeck H, Kvist L J and Landgren K ;. (2018). Gaining hope and self-confidence-An interview study of women's experience of treatment by art therapy for severe fear of childbirth. *Women & Birth: Journal of the Australian College of Midwives*, 31(4), pp.299-306.
- Wajid A, van Zanten and S V ; Mughal M K; Biringer A ; Austin M P; Vermeyden L ; Kingston D ;. (2019). Adversity in childhood and depression in pregnancy. *Archives of Women's Mental Health*, 23, pp.23.
- Walton G D, Ross L E; Stewart D E; Grigoriadis S and Dennis C L; Vigod S ;. (2014). Decisional conflict among women considering antidepressant medication use in pregnancy. *Archives of Women's Mental Health*, 17(6), pp.493-501.

Warriner S, Crane C and Dymond M ; Krusche A ;. (2018). An evaluation of mindfulness-based childbirth and parenting courses for pregnant women and prospective fathers/partners within the UK NHS (MBCP-4-NHS). *Midwifery*, 64, pp.1-10.

Webster J, Linnane J and Roberts J ; Starrenburg S ; Hinson J ; Dibley L ;. (2003). IDentify, Educate and Alert (IDEA) trial: an intervention to reduce postnatal depression. *BJOG: An International Journal of Obstetrics & Gynaecology*, 110(9), pp.842-6.

Weinreb L, Byatt N and Moore Simas ; T A ; Tenner K ; Savageau J A;. (2014). What happens to mental health treatment during pregnancy? Women's experience with prescribing providers. *Psychiatric Quarterly*, 85(3), pp.349-55.

Weinreb L, Wenz-Gross M and Upshur C ;. (2018). Postpartum outcomes of a pilot prenatal care-based psychosocial intervention for PTSD during pregnancy. *Archives of Women's Mental Health*, 21(3), pp.299-312.

Welch M G, Halperin M S; Austin J and Stark R I; Hofer M A; Hane A A; Myers M M;. (2016). Depression and anxiety symptoms of mothers of preterm infants are decreased at 4 months corrected age with Family Nurture Intervention in the NICU. *Archives of Women's Mental Health*, 19(1), pp.51-61.

Werner E A, Gustafsson H C; Lee S and Feng T ; Jiang N ; Desai P ; Monk C ;. (2016). PREPP: postpartum depression prevention through the mother-infant dyad. *Archives of Women's Mental Health*, 19(2), pp.229-42.

Wheatley Sandra L and Brugha T. (1999). 'Just because I like it doesn't mean it has to work': Personal experiences of an antenatal psychosocial intervention designed to prevent postnatal depression. *International Journal of Mental Health Promotion*, 1(1), pp.26-31.

White Jane, Thompson Lucy and Puckering Christine ; Waugh Harriet ; Henderson Marion ; MacBeth Angus ; Wilson Philip ;. (2015). Antenatal parenting support for vulnerable women. *British Journal of Midwifery*, 23(10), pp.724-732.

Wickberg B, Tjus T and Hwang P ;. (2005). Using the EPDS in routine antenatal care in Sweden: A naturalistic study. *Journal of Reproductive and Infant Psychology*, 23(1), pp.33-41.

Williams Nathalie B. (2008). Missouri Ozark women's experiences of living with postpartum depression symptoms. *Dissertation Abstracts International: Section B: The Sciences and Engineering*, 68(11-B), pp.7257.

Woolhouse H, Mercuri K and Judd F ; Brown S J;. (2014). Antenatal mindfulness intervention to reduce depression, anxiety and stress: a pilot randomised controlled trial of the MindBabyBody program in an Australian tertiary maternity hospital. *BMC Pregnancy & Childbirth*, 14, pp.369.

Wright B, Tindall L and Littlewood E ; Allgar V ; Abeles P ; Trepel D ; Ali S ;. (2017). Computerised cognitive-behavioural therapy for depression in adolescents: feasibility results and 4-month outcomes of a UK randomised controlled trial. *BMJ Open*, 7(1), pp.e012834.

Wroblewski M and Tallon D . (2004). Implementing a comprehensive postpartum depression support program. *AWHONN Lifelines*, 8(3), pp.248-52.

Wu Wan-Ru and Hung Chich-Hsiu . (2019). Impact of a peer virtual community on pregnant women's well-being: A repeated-measure and quasi-experimental study. *Journal of Advanced Nursing (John Wiley & Sons and Inc.)*, 75(5), pp.1099-1107.

Yaman S and Altay N . (2015). Posttraumatic stress and experiences of parents with a newborn in the neonatal intensive care unit. *Journal of Reproductive and Infant Psychology*, 33(2), pp.140-152.

Yawn B P, Pace W and Wollan P C; Bertram S ; Kurland M ; Graham D ; Dietrich A ;. (2009). Concordance of Edinburgh Postnatal Depression Scale (EPDS) and Patient Health Questionnaire (PHQ-9) to assess increased risk of depression among postpartum women. *Journal of the American Board of Family Medicine: JABFM*, 22(5), pp.483-91.

Yawn B P, Pace W and Dietrich A ; Bertram S ; Kurland M ; Graham D ; Huff J ; Rocca L ; Wollan P ;. (2010). Practice benefit from participating in a practice-based research network study of postpartum depression: a national research network (NRN) report. *Journal of the American Board of Family Medicine: JABFM*, 23(4), pp.455-64.

Yawn B P, Dietrich A J; Wollan P and Bertram S ; Graham D ; Huff J ; Kurland M ; Madison S ; Pace W D; practices Trippd ;. (2012). TRIPPD: a practice-based network effectiveness study of postpartum depression screening and management. *Annals of Family Medicine*, 10(4), pp.320-9.

Yazdanimehr R, Omid A and Sadat Z ; Akbari H ;. (2016). The Effect of Mindfulness-integrated Cognitive Behavior Therapy on Depression and Anxiety among Pregnant Women: a Randomized Clinical Trial. *Journal of Caring Sciences*, 5(3), pp.195-204.

Ying L and Feng C . (2017). The occurrence of depression and nursing intervention for mothers of preterm infants. *Biomedical Research (India)*, 28(12), pp.5530-5534.

Yonkers K A, Smith M V; Lin H and Howell H B; Shao L ; Rosenheck R A;. (2009). Depression screening of perinatal women: an evaluation of the healthy start depression initiative. *Psychiatric Services*, 60(3), pp.322-8.

Zayas L H and McKee M D; Jankowski K R;. (2004). Adapting psychosocial intervention research to urban primary care environments: a case example. *Annals of Family Medicine*, 2(5), pp.504-8.

Zhang Huaiyu and Emory Eugene K. (2015). A mindfulness-based intervention for pregnant African-American women. *Mindfulness*, 6(3), pp.663-674.

Zhang Yanjun and Jin Shenghua . (2016). The impact of social support on postpartum depression: The mediator role of self-efficacy. *Journal of Health Psychology*, 21(5), pp.720-726.

Zhao Y, Munro-Kramer M L and Shi S ; Wang J ; Zhao Q ;. (2019). Effects of antenatal depression screening and intervention among Chinese high-risk pregnant women with medically defined complications: A randomized controlled trial. *Early intervention in psychiatry*, 13(5), pp.1090-1098.

Zlotnick C, Capezza N M and Parker D ;. (2011). An interpersonally based intervention for low-income pregnant women with intimate partner violence: a pilot study. *Archives of Women's Mental Health*, 14(1), pp.55-65.

Zlotnick Caron, Wernette Golfo Tzilos and Raker Christina A;. (2018). A randomized controlled trial of a computer-based brief intervention for victimized perinatal women seeking mental health treatment. *Archives of Women's Mental Health*, , pp.No Pagination Specified.

Zourladani Athanasia, Tsaloglidou Areti and Tzetzis George ; Tsorbatzoudis Charalambos ; Matziari Chrysoula ;. (2011). The effect of a low impact exercise training programme on the well-being of Greek postpartum women: A randomised controlled trial. *International SportMed Journal*, 12(1), pp.30-38.

### **Not primary**

Andriotti T, Stavale R and Nafee T ; Fakhry S ; Mohamed M M. A; Sofiyeva N ; Ganho-Avila A ; Bogner A ; Barbosa S P; Piton L S; Hirayama A L. S; Gaccia G ; Smith-Howard Junior ; T P ; Miranda P C; Reyes K J. C; Gragera A ; Nishiwaki H ; Boechat-Barros R ;. (2017). ASSERT trial - How to assess the safety and efficacy of a high frequency rTMS in postpartum depression ? A multicenter, double blinded, randomized, placebo-controlled clinical trial. *Contemporary Clinical Trials Communications*, 5, pp.86-91.

Battle Cynthia L, Uebelacker Lisa A; Howard Margaret and Castaneda Michelle ;. (2010). Prenatal yoga and depression during pregnancy. *Birth: Issues in Perinatal Care*, 37(4), pp.353-354.

Battle C L and Abrantes A M; Schofield C A; Kraines M A;. (2015). Physical Activity as an Intervention for Antenatal Depression: Rationale for Developing Tailored Exercise Programs for Pregnant Women with Depression. *Journal of Midwifery & Women's Health*, 60(5), pp.479-82.

Baumgartner J N, Kaaya S and Siril H ;. (2015). Mental health screenings for couples at churches in Nigeria: a strategy for enhancing community-based maternal mental health services in low-resource settings. *Social Psychiatry & Psychiatric Epidemiology*, 50(3), pp.497-9.

Beck C T. (1995). Screening methods for postpartum depression. *JOGNN - Journal of Obstetric, Gynecologic and Neonatal Nursing*, 24(4), pp.308-12.

Boyce Philip M and Judd Fiona. (2019). Screening for perinatal depression: is it enough?. *Medical Journal of Australia*, 210(1), pp.19-20.

Buist A E, Barnett B E; Milgrom J and Pope S ; Condon J T; Ellwood D A; Boyce P M; Austin M P; Hayes B A;. (2002). To screen or not to screen--that is the question in perinatal depression. *Medical Journal of Australia*, 177(S7), pp.S101-5.

Buist A. (2002). Mental health in pregnancy: The sleeping giant. *Australasian Psychiatry*, 10(3), pp.203-206.

Canady Valerie A. (2018). Coordinated, community support drives effort to address postpartum depression. *Mental Health Weekly*, 28(28), pp.1-7.

Caramlau I, Barlow J and Sembi S ; McKenzie-McHarg K ; McCabe C ;. (2011). Mums 4 Mums: structured telephone peer-support for women experiencing postnatal depression. Pilot and exploratory RCT of its clinical and cost effectiveness. *Trials [Electronic Resource]*, 12, pp.88.

Dennis C. (2006). Counselling and cognitive behavioural therapy reduce anxiety and depression in women with postnatal depression. *Evidence Based Mental Health*, 9(2), pp.50-50.

Felder J N. (2019). Implementing the USPSTF Recommendations on Prevention of Perinatal Depression-Opportunities and Challenges. *JAMA Internal Medicine*, 179(4), pp.467-468.

Fonte Julia and Horton-Deutsch Sara . (2005). Treating Postpartum Depression in Immigrant Muslim Women. *Journal of the American Psychiatric Nurses Association*, 11(1), pp.39-44.

Freeman M P. (2019). Perinatal Depression: Recommendations for Prevention and the Challenges of Implementation. *JAMA*, 321(6), pp.550-552.

Haas Iii and William C . (2016). Yoga for Prenatal Depression. *Integrative Medicine Alert*, 19(2), pp.22-23.

Haynes E. (2018). Reaching women with perinatal mental illness at the booking-in appointment. *International Journal of Health Governance*, 23(1), pp.38-45.

Holden J M. (1991). Postnatal depression: its nature, effects, and identification using the Edinburgh Postnatal Depression scale. *Birth*, 18(4), pp.211-21.

Hughes L P, Austin-Ketch T and Volpe E M; Campbell-Heider N ;. (2017). Identification and Treatment of Adolescents With Perinatal Anxiety and Depression. *Journal of Psychosocial Nursing & Mental Health Services*, 55(6), pp.23-29.

Hynie Michaela, Umubyeyi Benoitte and Gasanganwa Marie Claire; Bohr Yvonne ; McGrath Susan ; Umuziga Providence ; Mukamsanga Beata ;. (2015). Community resilience and community interventions for post-natal depression: Reflecting on maternal mental health in Rwanda. In: , ed., *Women's mental health: Resistance and resilience in community and society*. Cham, Switzerland: Springer International Publishing; Switzerland, pp.343-356.

Keefe R H, Brownstein-Evans C and Rouland Polmanteer ; R S ;. (2016). Addressing access barriers to services for mothers at risk for perinatal mood disorders: A social work perspective. *Social Work in Health Care*, 55(1), pp.1-11.

Kenyon S, Jolly K and Hemming K ; Ingram L ; Gale N ; Dann S A; Chambers J ; MacArthur C ;. (2012). Evaluation of Lay Support in Pregnant women with Social risk (ELSIPS): a randomised controlled trial. *BMC Pregnancy & Childbirth*, 12, pp.11.

Lenz B, Eichler A and Schwenke E ; Buchholz V N; Hartwig C ; Moll G H; Reich K ; Muhle C ; Volz B ; Titzmann A ; Beckmann M W; Heinrich H ; Kornhuber J ; Fasching P A;. (2018). Mindfulness-based Stress Reduction in Pregnancy: An App-Based Programme to Improve the Health of Mothers and Children (MINDFUL/PMI Study). *Geburtshilfe und Frauenheilkunde*, 78(12), pp.1283-1291.

Logsdon M C and Gennaro S. (2005). Bioecological model for guiding social support research and interventions with pregnant adolescents. *Issues in Mental Health Nursing*, 26(3), pp.327-39.

Logsdon M C, Wisner K and Billings D M; Shanahan B ;. (2006). Raising the awareness of primary care providers about postpartum depression. *Issues in Mental Health Nursing*, 27(1), pp.59-73.

- Marcano Belisario, J S and Doherty K ; O'Donoghue J ; Ramchandani P ; Majeed A ; Doherty G ; Morrison C ; Car J ;. (2017). A bespoke mobile application for the longitudinal assessment of depression and mood during pregnancy: protocol of a feasibility study. *BMJ Open*, 7(5), pp.e014469.
- Milgrom J, Mendelsohn J and Gemmill A W;. (2011). Does postnatal depression screening work? Throwing out the bathwater, keeping the baby. *Journal of Affective Disorders*, 132(3), pp.301-10.
- Milgrom J and Gemmill A . (2014). Feasibility and efficacy of an internet treatment for postnatal depression utilising a behavioural activation approach. *Evidence-Based Nursing*, 17(4), pp.102.
- Pedrina F. (2004). Group Therapy with Mothers and Babies in Postpartum Crises: Preliminary Evaluation of A Pilot Project. *Group Analysis*, 37(1), pp.137-151.
- Posmontier B. (2010). The role of midwives in facilitating recovery in postpartum psychosis. *Journal of Midwifery & Women's Health*, 55(5), pp.430-7.
- Sharma V, Al-Farayedhi M and Doobay M ; Baczynski C ;. (2018). Should all women with postpartum depression be screened for bipolar disorder?. *Medical Hypotheses*, 118, pp.26-28.
- Shorey S, Chan W C and Chong Y S; He H G;. (2015). A randomized controlled trial of the effectiveness of a postnatal psychoeducation programme on outcomes of primiparas: study protocol. *Journal of Advanced Nursing*, 71(1), pp.193-203.
- Silverstein Michael, Feinberg Emily and Hegel Mark ;. (2011). Preventing depression of mothers of preterm infants. *Psychiatric Services*, 62(1), pp.101.
- Sit D K, Flint C and Svidergol D ; White J ; Wimer M ; Bish B ; Wisner K L;. (2009). Best practices: an emerging best practice model for perinatal depression care. *Psychiatric Services*, 60(11), pp.1429-31.
- Smith D M, Whitworth M and Sibley C ; Taylor W ; Gething J ; Chmiel C ; Lavender T ;. (2010). The design of a community lifestyle programme to improve the physical and psychological well-being of pregnant women with a BMI of 30 kg/m<sup>2</sup> or more. *BMC Public Health*, 10, pp.284.
- SmithBattle L and Freed P . (2016). Teen Mothers' Mental Health. *MCN and American Journal of Maternal Child Nursing*, 41(1), pp.31-6; quiz E3-4.
- Springate B and Chaudron L . (2006). Mental health providers' self-reported expertise and treatment of perinatal depression. *Archives of Women's Mental Health*, 9(1), pp.60-61.
- Terrazas C, Segre L S and Wolfe C ;. (2018). Moving beyond depression screening: integrating perinatal depression treatment into OB/GYN practices. *Primary Health Care Research & Development*, , pp.1-9.
- Yawn B P, Olson A L; Bertram S and Pace W ; Wollan P ; Dietrich A J;. (2012). Postpartum Depression: Screening, Diagnosis, and Management Programs 2000 through 2010. *Depression Research and Treatment*, 2012, pp.363964.

Zelkowitz P, Feeley N and Shrier I ; Stremler R ; Westreich R ; Dunkley D ; Steele R ; Rosberger Z ; Lefebvre F ; Papageorgiou A ;. (2009). The Cues and Care trial: a randomized controlled trial of an intervention to reduce maternal anxiety and improve developmental outcomes in very low birthweight infants. *Neonatal Intensive Care*, 22(1), pp.31-36.

## **Review 2:**

### **Can't find full text**

Byatt N, Xu W and Levin L L; Moore Simas ; T A ;. (2019). Perinatal depression care pathway for obstetric settings. *International Review of Psychiatry*, 31(3), pp.210-228.

Cannon C and Nasrallah H A. (2019). A focus on postpartum depression among African American women: A literature review. *Annals of Clinical Psychiatry*, 31(2), pp.138-143.

Doucet Shelley Anne. (2013). Postpartum psychosis: Support needs of mothers and fathers. *Dissertation Abstracts International: Section B: The Sciences and Engineering*, 73(11-B(E)), pp.No Pagination Specified.

Hippman C and Balneaves L G. (2018). Women's decision making about antidepressant use during pregnancy: A narrative review. *Depression and Anxiety*, 35(12), pp.1158-1167.

Mason Malgorzata. (2020). POSTNATAL DEPRESSION: HV SUPPORT EXAMINED. *Community Practitioner*, 93(3), pp.44-47.

### **Not mental health**

Perez A, Panagiotopoulou E and Curtis P ; Roberts R ;. (2021). Barriers and facilitators to mood and confidence in pregnancy and early parenthood during COVID-19 in the UK: mixed-methods synthesis survey. *BJPsych Open*, 7(4), pp.e107.

Santiago Mda, C and Figueiredo M H;. (2015). Immigrant women's perspective on prenatal and postpartum care: systematic review. *Journal of Immigrant & Minority Health*, 17(1), pp.276-84.

Sivertsen N, Anikeeva O and Deverix J ; Grant J ;. (2020). Aboriginal and Torres Strait Islander family access to continuity of health care services in the first 1000 days of life: a systematic review of the literature. *BMC Health Services Research*, 20(1), pp.829.

### **Not about access to PMH care**

Ammerman R T, Putnam F W; Bosse N R; Teeters A R; Van Ginkel and J B ;. (2010). Maternal depression in home visitation: a systematic review. *Aggression & Violent Behavior*, 15(3), pp.191-200.

Brandao T, Brites R and Nunes O ; Hipolito J ;. (2020). Posttraumatic Growth After Childbirth in Women: A Systematic Review. *Journal of Clinical Psychology in Medical Settings*, 27(2), pp.318-330.

Bright K S, Norris J M; Letourneau N L; King Rosario and M ; Premji S S;. (2018). Prenatal Maternal Anxiety in South Asia: A Rapid Best-Fit Framework Synthesis. *Frontiers in psychiatry Frontiers Research Foundation*, 9, pp.467.

Broughton L C and Medicott N J; Smith A J;. (2021). Effectiveness of patient decision aids in women considering psychotropic medication use during pregnancy: a literature review. *Archives of Women's Mental Health*, 24(4), pp.569-578.

Dixon S and Dantas J A. (2017). Best practice for community-based management of postnatal depression in developing countries: A systematic review. *Health Care for Women International*, 38(2), pp.118-143.

Fagbemi Folashade. (2019). Impact of social support on postnatal depression among adolescent mothers: a systematic literature review. *MIDIRS Midwifery Digest*, 29(1), pp.20-26.

Fonseca A, Ganho-Avila A , Lambregtse-van den Berg and M; Lupattelli A ; Rodriguez-Munoz M F; Ferreira P ; Rados S N; Bina R ;. (2020). Emerging issues and questions on peripartum depression prevention, diagnosis and treatment: a consensus report from the cost action riseup-PPD. *Journal of Affective Disorders*, 274, pp.167-173.

Gillis Brenna D and Parish Abby L;. (2019). Group-based interventions for postpartum depression: An integrative review and conceptual model. *Archives of Psychiatric Nursing*, 33(3), pp.290-298.

Kita Sachiko, Minatani Mariko and Hikita Naoko ; Matsuzaki Masayo ; Shiraishi Mie ; Haruna Megumi ;. (2015). A Systematic Review of the Physical, Mental, Social, and Economic Problems of Immigrant Women in the Perinatal Period in Japan. *Journal of Immigrant & Minority Health*, 17(6), pp.1863-1881.

Langan R C and Goodbred A J;. (2016). Identification and management of peripartum depression. *American Family Physician*, 93(10), pp.852-858.

Lara-Cinisomo S, Ramirez Olarte and A ; Rosales M ; Barrera A Z;. (2021). A Systematic Review of Technology-Based Prevention and Treatment Interventions for Perinatal Depression and Anxiety in Latina and African American Women. *Maternal & Child Health Journal*, 25(2), pp.268-281.

Lau Y, Htun T P and Wong S N; Tam W S. W; Klainin-Yobas P ;. (2017). Therapist-Supported Internet-Based Cognitive Behavior Therapy for Stress, Anxiety, and Depressive Symptoms Among Postpartum Women: A Systematic Review and Meta-Analysis. *Journal of Medical Internet Research*, 19(4), pp.e138.

Leahy-Warren P and McCarthy G . (2007). Postnatal depression: prevalence, mothers' perspectives, and treatments. *Archives of Psychiatric Nursing*, 21(2), pp.91-100.

Lee D T. S and Chung T K. H;. (2007). Postnatal depression: an update. *Best Practice and Research in Clinical Obstetrics and Gynaecology*, 21(2), pp.183-191.

Lee E W, Denison F C; Hor K and Reynolds R M;. (2016). Web-based interventions for prevention and treatment of perinatal mood disorders: a systematic review. *BMC Pregnancy & Childbirth*, 16, pp.38.

McGeough Christine, Walsh Aisling and Clyne Barbara ;. (2020). Barriers and facilitators perceived by women while homeless and pregnant in accessing antenatal and or postnatal healthcare: A qualitative evidence synthesis. *Health & Social Care in the Community*, 28(5), pp.1380-1393.



Patterson J, Hollins Martin and C ; Karatzias T ;. (2019). PTSD post-childbirth: a systematic review of women's and midwives' subjective experiences of care provider interaction. *Journal of Reproductive & Infant Psychology*, 37(1), pp.56-83.

Pettman D, O'Mahen H and Skoog Svanberg ; A ; von Essen ; L ; Axfors C ; Blomberg O ; Woodford J ;. (2019). Effectiveness and acceptability of cognitive-behavioural therapy based interventions for maternal peripartum depression: a systematic review, meta-analysis and thematic synthesis protocol. *BMJ Open*, 9(12), pp.e032659.

Prady S L, Endacott C and Dickerson J ; Bywater T J; Blower S L;. (2021). Inequalities in the identification and management of common mental disorders in the perinatal period: An equity focused re-analysis of a systematic review. *PLoS ONE [Electronic Resource]*, 16(3), pp.e0248631.

Qutteina Y, Nasrallah C and James-Hawkins L ; Nur A A; Yount K M; Hennink M ; Abdul Rahim ; H F ;. (2018). Social resources and Arab women's perinatal mental health: A systematic review. *Women & Birth: Journal of the Australian College of Midwives*, 31(5), pp.386-397.

Rayment-Jones H, Harris J and Harden A ; Khan Z ; Sandall J ;. (2019). How do women with social risk factors experience United Kingdom maternity care? A realist synthesis. *Birth*, 46(3), pp.461-474.

Shorey S and Wong P Z. E. (2020). Traumatic Childbirth Experiences of New Parents: A Meta-Synthesis. *Trauma Violence & Abuse*, , pp.1524838020977161.

Suwalska J, Napierala M and Bogdanski P ; Lojko D ; Wszolek K ; Suchowiak S ; Suwalska A ;. (2021). Perinatal mental health during covid-19 pandemic: An integrative review and implications for clinical practice. *Journal of Clinical Medicine*, 10(11), pp..

Syka Andria. (2015). REVIEW PAPER. Depression in Pregnancy and Ways of Dealing. *International Journal of Caring Sciences*, 8(1), pp.231-237.

Weiss-Laxer N S, Platt R and Osborne L M; Kimmel M ; Solomon B S; Mendelson T ; Webb L ; Riley A W;. (2016). Beyond screening: A review of pediatric primary care models to address maternal depression. *Pediatric Research*, 79(1-2), pp.197-204.

Wigert Helena, Nilsson Christina and Dencker Anna ; Begley Cecily ; Jangsten Elisabeth ; Sparud-Lundin Carina ; Mollberg Margareta ; Patel Harshida ;. (2020). Women's experiences of fear of childbirth: a metasynthesis of qualitative studies. *International Journal of Qualitative Studies on Health & Well-Being*, 15(1), pp.1-11.

Wittkowski A, Gardner P L and Bunton P ; Edge D ;. (2014). Culturally determined risk factors for postnatal depression in Sub-Saharan Africa: a mixed method systematic review. *Journal of Affective Disorders*, 163, pp.115-24.

Wong J and Fisher J . (2009). The role of traditional confinement practices in determining postpartum depression in women in Chinese cultures: a systematic review of the English language evidence. *Journal of Affective Disorders*, 116(3), pp.161-9.

Yargawa Judith and Leonardi-Bee Jo . (2015). Male involvement and maternal health outcomes: systematic review and meta-analysis. *Journal of Epidemiology & Community Health*, 69(6), pp.604-612.

### **Barriers not from mothers perspective**

Abdollahi F, Lye M S and Zain A M; Ghazali S S; Zarghami M ;. (2011). Postnatal depression and its associated factors in women from different cultures. *Iranian Journal of Psychiatry and Behavioral Sciences*, 5(2), pp.5-11.

Arefadib N, Cooklin A and Nicholson J ; Shafiei T ;. (2021). Postnatal depression and anxiety screening and management by maternal and child health nurses in community settings: A scoping review. *Midwifery*, 100, pp.103039.

Coates D and Foureur M . (2019). The role and competence of midwives in supporting women with mental health concerns during the perinatal period: A scoping review. *Health & Social Care in the Community*, 27(4), pp.e389-e405.

Coleman V H, Morgan M A; Zinberg S and Schulkin J ;. (2006). Clinical approach to mental health issues among obstetrician-gynecologists: a review. *Obstetrical & Gynecological Survey*, 61(1), pp.51-8.

Long M M, Cramer R J; Jenkins J and Bennington L ; Paulson J F;. (2019). A systematic review of interventions for healthcare professionals to improve screening and referral for perinatal mood and anxiety disorders. *Archives of Women's Mental Health*, 22(1), pp.25-36.

Moore Simas, T A and Flynn M P; Kroll-Desrosiers A R; Carvalho S M; Levin L L; Biebel K ; Byatt N ;. (2018). A Systematic Review of Integrated Care Interventions Addressing Perinatal Depression Care in Ambulatory Obstetric Care Settings. *Clinical Obstetrics & Gynecology*, 61(3), pp.573-590.

Murthy S, Haeusslein L and Bent S ; Fitelson E ; Franck L S; Mangurian C ;. (2021). Feasibility of universal screening for postpartum mood and anxiety disorders among caregivers of infants hospitalized in NICUs: a systematic review. *Journal of Perinatology*, 10, pp.10.

Noonan M, Galvin R and Doody O ; Jomeen J ;. (2017). A qualitative meta-synthesis: public health nurses role in the identification and management of perinatal mental health problems. *Journal of Advanced Nursing*, 73(3), pp.545-557.

Wadephul F, Jarrett P M and Jomeen J ; Martin C R;. (2018). A mixed methods review to develop and confirm a framework for assessing midwifery practice in perinatal mental health. *Journal of Advanced Nursing*, 74(10), pp.2258-2272.

Wyatt Caroline. (2021). Postnatal mental distress: Exploring the experiences of professionals, mothers, and significant others. *Dissertation Abstracts International: Section B: The Sciences and Engineering*, 82(6-B), pp.No Pagination Specified.

### **Does not provide reasons for lack of access/no list of barriers/facilitators**

Byatt N, Levin L L and Ziedonis D ; Moore Simas ; T A ; Allison J ;. (2015). Enhancing Participation in Depression Care in Outpatient Perinatal Care Settings: A Systematic Review. *Obstetrics & Gynecology*, 126(5), pp.1048-58.

Chowdhary N, Sikander S and Atif N ; Singh N ; Ahmad I ; Fuhr D C; Rahman A ; Patel V ;. (2014). The content and delivery of psychological interventions for perinatal depression by

non-specialist health workers in low and middle income countries: a systematic review. *Best Practice & Research in Clinical Obstetrics & Gynaecology*, 28(1), pp.113-33.

des Anges , C and Asante-Ackuayi L ; Brown S ; Cardenas E ; Lee D ;. (2012). Evidence-Based Guidance for Culturally Sensitive Assessment and Interventions for Perinatal Depression in Black American Women: A Synthesis of Published Research, 2008-2011. *Journal of Primary Care and Community Health*, 3(4), pp.278-284.

Downe S M, Butler E and Hinder S ;. (2007). Screening tools for depressed mood after childbirth in UK-based South Asian women: a systematic review. *Journal of Advanced Nursing*, 57(6), pp.565-83.

El-Den S, O'Reilly C L and Chen T F;. (2015). A systematic review on the acceptability of perinatal depression screening. *Journal of Affective Disorders*, 188, pp.284-303.

Gearing R E, Alonzo D and Marinelli C ;. (2012). Maternal schizophrenia: psychosocial treatment for mothers and their children. *Clinical Schizophrenia & Related Psychoses*, 6(1), pp.27-33.

Gillham R and Wittkowski A . (2015). Outcomes for women admitted to a mother and baby unit: a systematic review. *International Journal of Women's Health*, 7, pp.459-76.

Lavender Theresa J, Ebert Lyn and Jones Donovan ;. (2016). An evaluation of perinatal mental health interventions: An integrative literature review. *Women & Birth*, 29(5), pp.399-406.

### **Perinatal mental health barriers to access not described separately to other services**

Breckenridge J P, Devaney J and Kroll T ; Lazenbatt A ; Taylor J ; Bradbury-Jones C ;. (2014). Access and utilisation of maternity care for disabled women who experience domestic abuse: a systematic review. *BMC Pregnancy & Childbirth*, 14, pp.234.

Rogers H J, Hogan L and Coates D ; Homer C S. E; Henry A ;. (2020). Responding to the health needs of women from migrant and refugee backgrounds-Models of maternity and postpartum care in high-income countries: A systematic scoping review. *Health & Social Care in the Community*, 30, pp.30.

### **No PRISMA search strategy**

Accortt E E and Wong M S;. (2017). It Is Time for Routine Screening for Perinatal Mood and Anxiety Disorders in Obstetrics and Gynecology Settings. *Obstetrical and Gynecological Survey*, 72(9), pp.553-568.

Austin M P and Priest S R;. (2005). Clinical issues in perinatal mental health: new developments in the detection and treatment of perinatal mood and anxiety disorders. *Acta Psychiatrica Scandinavica*, 112(2), pp.97-104.

Buist A E, Barnett B E. W; Milgrom J and Pope S ; Condon J T; Ellwood D A; Boyce P M; Austin M P. V; Hayes B A;. (2002). To screen or not to screen - That is the question in perinatal depression. *Medical Journal of Australia*, 177(7 SUPPL.), pp.S101-S105.

Byatt N, Simas T A and Lundquist R S; Johnson J V; Ziedonis D M;. (2012). Strategies for improving perinatal depression treatment in North American outpatient obstetric settings. *Journal of Psychosomatic Obstetrics & Gynecology*, 33(4), pp.143-61.

Gjerdingen D K and Yawn B P;. (2007). Postpartum depression screening: importance, methods, barriers, and recommendations for practice. *Journal of the American Board of Family Medicine: JABFM*, 20(3), pp.280-8.

Jarrett Patricia M. (2016). Pregnant women's experience of depression care. *Journal of Mental Health Training and Education & Practice*, 11(1), pp.33-47.

Johnson Samantha, Adam Simon and McIntosh Michele ;. (2020). The Lived Experience of Postpartum Depression: A Review of the Literature. *Issues in Mental Health Nursing*, 41(7), pp.584-591.

Liberto T L. (2012). Screening for depression and help-seeking in postpartum women during well-baby pediatric visits: an integrated review. *Journal of Pediatric Health Care*, 26(2), pp.109-17.

McLoughlin Jennifer. (2013). Stigma associated with postnatal depression: A literature review. *British Journal of Midwifery*, 21(11), pp.784-791.

O'Mahony J and Donnelly T . (2010). Immigrant and refugee women's post-partum depression help-seeking experiences and access to care: a review and analysis of the literature. *Journal of Psychiatric & Mental Health Nursing*, 17(10), pp.917-28.