

## **Supplementary Material 12: Examples of additional outcomes included:**

**1. Extended Activities of Daily Living.** Measured by scales such as Frenchay Activities Index <sup>1</sup>, Nottingham Extended Activities of Daily Living scale <sup>2</sup>, Lawton Instrumental Activities of Daily Living <sup>3</sup>, Rivermead Activities of Daily Living score <sup>4</sup>.

### **2. Quality of Life.**

- **Quality of Life** e.g. EQ5D <sup>5</sup>, Health- related quality of life scale <sup>6</sup>, Quality of Well Being Scale <sup>7, 8</sup>, SF36 <sup>9</sup>, Stroke Impact Scale <sup>10</sup>.
- **Social activities and participation** e.g. Australian Community Participation Questionnaire <sup>11</sup>, ICEpop CAAbility measure for Older people <sup>12</sup>,
- **Mobility, navigation and safety** e.g. Rivermead Mobility Index <sup>13</sup>, 6-minute walk test <sup>14</sup>, functional ambulation <sup>15</sup>, timed up and go <sup>16</sup>

### **3. Psychological and mental health**

- **Stroke survivors.** e.g. Hospital Anxiety and Depression Scale <sup>17</sup>, Beck Depression Inventory <sup>18</sup>, General Health Questionnaire <sup>19</sup>, Geriatric Depression Scale <sup>20</sup>, discharge destination.
- **Family, friends and carers.** E.g. The Carer Strain Index <sup>21</sup>, Perceived Caregiver Burden <sup>22</sup>.

**4. Perceptual function** e.g. Rivermead Perceptual Assessment Battery <sup>23</sup>, Motor Free Visual Perception <sup>24</sup>, Birmingham Object Recognition Battery <sup>25</sup>, Chessington Occupational Therapy <sup>26</sup>, Neurological Assessment Battery <sup>27</sup>.

**5. Adverse events** e.g. falls, death, fatigue, accident rates.

### ***Additional outcomes of Interest***

These were

- Sensation e.g. Visual acuity, visual fields,
- Motor ability (including balance) e.g. grip strength <sup>28</sup>, Berg Balance Scale <sup>29</sup>, Postural Assessment Scale for Stroke <sup>30</sup>, Motor Assessment Scale <sup>31</sup>, Fugl-Meyer Assessment <sup>32</sup>, Motricity Index <sup>33</sup>
- Cognition (including attention) e.g. Mini–Mental State Examination <sup>34</sup>, Wechsler Adult Intelligence Scale (WAIS) <sup>35</sup>, line bisection <sup>36</sup>, star cancellation <sup>37</sup>

The use of these outcomes was noted, as they were considered of potential interest to readers, but no further action taken.

## References

1. Holbrook M, Skilbeck CE. An activities index for use with stroke patients. *Age Ageing* 1983;12:166-70. <http://dx.doi.org/10.1093/ageing/12.2.166>
2. Nouri FM, Lincoln NB. An extended activities of daily living scale for stroke patients. *Clin Rehabil* 1987;1:301-5.
3. Graf C. The Lawton instrumental activities of daily living scale. *Am J Nurs* 2008;108:52-62; quiz -3. <http://dx.doi.org/10.1097/01.NAJ.0000314810.46029.74>
4. Lincoln NB, Edmans JA. A re-validation of the Rivermead ADL scale for elderly patients with stroke. *Age Ageing* 1990;19:19-24. <http://dx.doi.org/10.1093/ageing/19.1.19>
5. Balestroni G, Bertolotti G. [EuroQol-5D (EQ-5D): an instrument for measuring quality of life]. *Monaldi Arch Chest Dis* 2012;78:155-9. <http://dx.doi.org/10.4081/monaldi.2012.121>
6. Centers for Disease Control and Prevention. *Measuring Healthy Days. Population assessment of health-related quality of life*. USA: US Department of Health and Human Sciences, Centers for Disease Control and Prevention; 202031 March 2022).
7. Kaplan RM, Bush JW. Health-related quality of life measurements for evaluation research and policy analysis. *Health Psychol* 1982;1:61.
8. Kaplan RM, Bush JW, Berry CC. The reliability, stability and generalizability of a health status index. American Statistical Society; Washington, DC, abstract no. 153.
9. McHorney CA, Ware JE, Jr., Lu JF, Sherbourne CD. The MOS 36-item Short-Form Health Survey (SF-36): III. Tests of data quality, scaling assumptions, and reliability across diverse patient groups. *Med Care* 1994;32:40-66. <http://dx.doi.org/10.1097/00005650-199401000-00004>
10. Duncan PW, Bode RK, Min Lai S, Perera S, Glycine Antagonist in Neuroprotection Americans I. Rasch analysis of a new stroke-specific outcome scale: the Stroke Impact Scale. *Arch Phys Med Rehabil* 2003;84:950-63. [http://dx.doi.org/10.1016/s0003-9993\(03\)00035-2](http://dx.doi.org/10.1016/s0003-9993(03)00035-2)
11. Berry HL, Rodgers B, Dear KB. Preliminary development and validation of an Australian community participation questionnaire: types of participation and associations with distress in a coastal community. *Soc Sci Med* 2007;64:1719-37. <http://dx.doi.org/10.1016/j.socscimed.2006.12.009>
12. Al-Janabi H, Flynn TN, Coast J. Development of a self-report measure of capability wellbeing for adults: the ICECAP-A. *Qual Life Res* 2012;21:167-76. <http://dx.doi.org/10.1007/s11136-011-9927-2>
13. Collen FM, Wade DT, Robb GF, Bradshaw CM. The Rivermead Mobility Index: a further development of the Rivermead Motor Assessment. *Int Disabil Stud* 1991;13:50-4. <http://dx.doi.org/10.3109/03790799109166684>
14. Butland RJ, Pang J, Gross ER, Woodcock AA, Geddes DM. Two-, six-, and 12-minute walking tests in respiratory disease. *Br Med J (Clin Res Ed)* 1982;284:1607-8. <http://dx.doi.org/10.1136/bmj.284.6329.1607>
15. Holden MK, Gill KM, Magliozzi MR, Nathan J, Piehl-Baker L. Clinical gait assessment in the neurologically impaired. Reliability and meaningfulness. *Phys Ther* 1984;64:35-40. <http://dx.doi.org/10.1093/ptj/64.1.35>

16. Podsiadlo D, Richardson S. The timed "Up & Go": a test of basic functional mobility for frail elderly persons. *J Am Geriatr Soc* 1991;39:142-8. <http://dx.doi.org/10.1111/j.1532-5415.1991.tb01616.x>
17. Zigmond AS, Snaith RP. The hospital anxiety and depression scale. *Acta Psychiatr Scand* 1983;67:361-70. <http://dx.doi.org/10.1111/j.1600-0447.1983.tb09716.x>
18. McDowell I, Newell C. *Measuring Health. A Guide to Rating Scales and Questionnaires*. 2nd edn. New York: Oxford University Press; 1996.
19. Goldberg DP, Hillier VF. A scaled version of the General Health Questionnaire. *Psychol Med* 1979;9:139-45. <http://dx.doi.org/10.1017/s0033291700021644>
20. Stiles PG, McGarrah JE. The Geriatric Depression Scale: A comprehensive review. *Journal of Clinical Geropsychology* 1998;4:89-109. <http://dx.doi.org/10.1037/t00930-000>
21. Robinson BC. Validation of a Caregiver Strain Index. *J Gerontol* 1983;38:344-8. <http://dx.doi.org/10.1093/geronj/38.3.344>
22. Stommel M, Given CW, Given B. Depression as an Overriding Variable Explaining Caregiver Burdens. *Journal of Aging and Health* 2016;2:81-102. <http://dx.doi.org/10.1177/089826439000200106>
23. Whiting S, Lincoln NB, Bhavnani G, Cockburn J. *The Rivermead Perceptual Assessment Battery*. Windsor: NFER-Nelson; 1985.
24. Colarusso RP, Hammill DD. *The Motor Free Visual Perception Test (MVPT-3)*. Navato, CA: Academic Therapy Publications; 2003.
25. Riddoch MJ, Humphreys GW. *Birmingham Object Recognition Battery*. Hove, UK: Lawrence Erlbaum Associates, Publishers; 1993.
26. Tyerman R, Tyeman A, Howard P, Hadfield C. *The Chessington OT Neurological Assessment Battery introduction manual*. Nottingham, UK: Nottingham Rehab Limited; 1986.
27. Laver AJ, Huchison S. The Performance and Experience of Normal Elderly People on the Chessington Occupational Therapy Neurological Assessment Battery (COTNAB). *British Journal of Occupational Therapy* 2016;57:137-42. <http://dx.doi.org/10.1177/030802269405700410>
28. Bohannon RW. Dynamometer measurements of hand-grip strength predict multiple outcomes. *Percept Mot Skills* 2001;93:323-8. <http://dx.doi.org/10.2466/pms.2001.93.2.323>
29. Berg KO, Wood-Dauphinee SL, Williams JI, Maki B. Measuring balance in the elderly: validation of an instrument. *Can J Public Health* 1992;83 Suppl 2:S7-11.
30. Benaim C, Perennou DA, Villy J, Rousseaux M, Pelissier JY. Validation of a standardized assessment of postural control in stroke patients: the Postural Assessment Scale for Stroke Patients (PASS). *Stroke* 1999;30:1862-8. <http://dx.doi.org/10.1161/01.str.30.9.1862>
31. Carr JH, Shepherd RB, Nordholm L, Lynne D. Investigation of a new motor assessment scale for stroke patients. *Phys Ther* 1985;65:175-80. <http://dx.doi.org/10.1093/ptj/65.2.175>
32. Fugl-Meyer AR. Post-stroke hemiplegia assessment of physical properties. *Scand J Rehabil Med Suppl* 1980;7:85-93.
33. Demeurisse G, Demol O, Robaye E. Motor evaluation in vascular hemiplegia. *Eur Neurol* 1980;19:382-9. <http://dx.doi.org/10.1159/000115178>
34. Folstein MF, Robins LN, Helzer JE. The Mini-Mental State Examination. *Arch Gen Psychiatry* 1983;40:812. <http://dx.doi.org/10.1001/archpsyc.1983.01790060110016>
35. Wechsler D. *Wechsler Adult Intelligence Scale--Fourth Edition (WAIS-IV)*. APA PsycNet; 2008. URL: <https://psycnet.apa.org/doiLanding?doi=10.1037%2Ft15169-000> (Accessed 25 March, 2022). <http://dx.doi.org/10.1037/t15169-000>

36. Plummer P, Morris ME, Dunai J. Assessment of unilateral neglect. *Phys Ther* 2003;83:732-40.
37. Wilson B, Cockburn J, Halligan P. Development of a behavioral test of visuospatial neglect. *Arch Phys Med Rehabil* 1987;68:98-102.