

Escalated dose of imatinib for patients with gastro intestinal stromal tumours

Assessor initials: Date:

Study identifier (Surname of first author + year of publication)

Type of study

Is the study an RCT in which all participants are randomised to imatinib, sunitinib or best supportive care (either provided in addition to imatinib or sunitinib or as only care)?




OR

Is the study a non-randomised comparative study on patients using either imatinib or sunitinib or best supportive care?

OR

Is the study case series or case study of more than one patient on same type of diagnosis?

Yes Unclear No

Go to next question **Exclude**




Participants in the study

Does the study contain participants with KIT (CD117) positive unresectable and/or metastatic malignant gastrointestinal stromal tumours (GIST)?

- Unresectable
 Metastatic

Does the study state that disease has progressed on treatment with imatinib at a dose of 400 mg/day? Yes No

Yes Unclear No

Go to next question **Exclude**

Doses and other comparisons

Does the study contain at least one group using escalated doses of imatinib (600mg or 800mg per day)?




OR

Does the study contain at least one group using sunitinib within its recommended dose range (i.e. 25-75 mg/day)?

OR

Does the study contain at least one group receiving best supportive care

Yes Unclear No




Go to next question **Exclude**

Outcomes reported

Does the study report any one of the following outcomes?

- Overall response
 Overall survival
 Disease-free survival
 Progression-free survival
 Time to treatment failure
 Health-related quality of life
 Adverse effects of treatment

Yes Unclear No

Go to next question **Exclude**

Decision

Include **Unclear** **Exclude**

Clarification
required