

| Reference | Contact method(s) |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------|
| Koefoed BG, Gullov AL, Pedersen TS, Petersen P. Dropout and withdrawal from warfarin and aspirin therapy in patients with atrial fibrillation. <i>Thromb Haemost</i> 1997;(Suppl.):83-4 | E-mail |
| Lavitola PL, Spina GS, Sampaio RO, Tarasoutchi F, Grinberg M. Bleeding during oral anticoagulant therapy: warning against a greater hazard. <i>Arq Bras Cardiol</i> 2009; 93 :174-9 | Post, e-mail |
| Levine MN, Raskob G, Hirsh J. Risk of haemorrhage associated with long term anticoagulant therapy. <i>Drugs</i> 1985; 30 :444-60 | Post, e-mail |
| Llobera J, Canameras N, Mas MA, Robles M, Llorach I, Miralles R, et al. [Atrial fibrillation and thromboembolic risk in the elderly.] <i>Rev Multidisciplin Gerontol</i> 2007; 17 :43-8 | Contact details not found |
| Matsuo S, Nakamura Y, Kinoshita M. Warfarin reduces silent cerebral infarction in elderly patients with atrial fibrillation. <i>Coron Artery Dis</i> 1998; 9 :223-6 | Post, e-mail |
| Neutel JM, Smith DHG. A randomised crossover study to compare the efficacy and tolerability of Barr Warfarin sodium to the currently available Coumadin. <i>Cardiovasc Rev Rep</i> 1998; 19 :49-59 | Post |
| Ortiz MR, Sanchez MA, Ortega MD, Rubio DM, Del Prado JMA, Zapata MF, et al. [Anticoagulation in patients aged less than 75 years with atrial fibrillation.] <i>Salud Cienc</i> 2008; 16 :164-7 | Post, e-mail |